

Erdős-Hajnal Sets and Semigroup Decompositions

Joshua N. Cooper ^{*}

Courant Institute of Mathematics, NYU

New York, NY

jcooper@cims.nyu.edu

September 14, 2004

Abstract

Define a set of lines in \mathbb{R}^3 to be “stacked” with respect to $\mathbf{v} \in \mathbb{R}^3$ if, from a vantage point far away in the direction of \mathbf{v} , the lines are linearly ordered by the “crossing over” relation. Given a collection of skew lines and a point \mathbf{v} , we ask, what is the largest stacked subset that must be present among the lines? This question, which appears in [6], is intimately related to the well-known Erdős-Hajnal conjecture via the Milnor-Thom theorem. It was recently resolved by a powerful and very general theorem of Alon, Pach, Pinchasi, Radoičić, and Sharir ([1]). We describe these results and discuss several related issues, including a generalization to “Erdős-Hajnal sets” and an intriguing problem concerning the decomposability of semi-algebraic sets: Do all semi-algebraic sets belong to the set algebra generated by semigroups in \mathbb{R}^d ? Our main result is a resolution of this question in dimensions 1 and 2.

Suppose we have a collection \mathcal{L} of n lines and a direction \mathbf{v} in \mathbb{R}^3 . Define T to be the set of the pairs $(\ell_1, \ell_2) \in \mathcal{L} \times \mathcal{L}$ so that ℓ_1 “crosses over” ℓ_2 from the perspective of a point very far away in the direction of \mathbf{v} . If the lines are pairwise skew, then T is a tournament. In [6], the authors ask for the size of its largest transitive subtournament: a set of lines which are linearly ordered by the “crossing over” relation, i.e., which appear to be “stacked” (q.v. Figure 1).

By Ramsey’s Theorem, every tournament, including T , must have a transitive subtournament of size $\Omega(\log n)$. Perhaps we can hope for more, however: T is very special, in that it is the result of a very specific construction. Surely, not all T can arise in this way from line configurations in \mathbb{R}^3 ...?

Indeed, the answer is “no”. There are very few such tournaments. To see this, we first make precise the notion of “crossing over”. Parameterize lines in \mathbb{R}^3 as follows. Consider the plane perpendicular to \mathbf{v} that passes through the origin. Call this the xy -plane. Then, consider the two planes $x = 1$ and $x = -1$. Each line ℓ crosses each plane in one point: $(1, a, b)$ and $(-1, c, d)$, respectively. (If ℓ is parallel to one of

^{*}Research supported by NSF Grant DMS-0303272.




Figure 1: Stacked lines vs. non-stacked lines.

them, we may rotate the xy -plane and reparameterize so this is no longer the case. Since n is finite, this can be done for all the lines simultaneously.) Therefore ℓ can be described uniquely by the 4-tuple (a, b, c, d) . Consider two lines, $\ell_1 = (a_1, b_1, c_1, d_1)$ and $\ell_2 = (a_2, b_2, c_2, d_2)$. They each project to a line on the xy -plane, and cross at a point (x, y) , so ℓ_1 contains a point (x, y, z_1) and ℓ_2 contains (x, y, z_2) . Whichever ℓ_i has the larger z_i “crosses over” the other. We may describe ℓ_i as

$$\{(1, a_i, b_i) + t_i(-2, c_i - a_i, d_i - b_i) : t_i \in \mathbb{R}\},$$

and its projection ℓ'_i onto the xy -plane as

$$\{(1, a_i) + t_i(-2, c_i - a_i) : t_i \in \mathbb{R}\}.$$

Then ℓ'_1 and ℓ'_2 intersect when $t_1 = t_2$ and $t_1(c_1 - a_1) + a_1 = t_1(c_2 - a_2) + a_2$, i.e., $t_1 = (a_1 - a_2)/((a_1 - a_2) - (c_1 - c_2))$. Therefore,

$$\begin{aligned} z_1 - z_2 &= b_1 - b_2 + (a_1 - a_2) \cdot \frac{(d_1 - d_2) - (b_1 - b_2)}{(a_1 - a_2) - (c_1 - c_2)} \\ &= \frac{(a_1 - a_2)(d_1 - d_2) - (b_1 - b_2)(c_1 - c_2)}{(a_1 - a_2) - (c_1 - c_2)}. \end{aligned}$$

Defining $g(a, b, c, d) = (a - c)(ad - bc)$, we have that ℓ_1 passes over ℓ_2 iff

$$g(a_1 - a_2, b_1 - b_2, c_1 - c_2, d_1 - d_2) > 0.$$

To show that this construction can give rise to only a small fraction of all possible tournaments, we employ a standard device: the Milnor-Thom Theorem. The following version of this important result appears in [8].

Theorem 1 (Milnor '64, Thom '65). *Let R be the number of ± 1 -vectors in the set*

$$\{(g_1(\mathbf{x}), \dots, g_m(\mathbf{x})) : \mathbf{x} \in \mathbb{R}^k\}$$

where $g_i = \text{sgn} \circ f_i$, $1 \leq i \leq m$, and each f_i is a polynomial of degree at most d in k variables. Then

$$R \leq \left(\frac{4edm}{k} \right)^k.$$

In this case, the tournament T is determined by the sign of the $m = \binom{n}{2}$ polynomials $g(a_i - a_j, b_i - b_j, c_i - c_j, d_i - d_j)$, $1 \leq i < j \leq n$, of degree $d = 3$ in the $k = 4n$ variables, so the number of such tournaments is at most

$$(3e(n-1)/2)^{4n} = 2^{O(n \log n)}.$$

On the other hand, the total number of tournaments on n vertices is $2^{\binom{n}{2}}$, so this is a vanishingly small fraction. In particular, there is some T_0 which can never be a subtournament of a tournament arising from a line configuration in \mathbb{R}^3 .

What does this say about the original question, i.e., what is the largest transitive subtournament of a tournament T arising from the crossing patterns of a set of n lines? Here, the Erdős-Hajnal conjecture plays a role. Define $\text{hom}(G)$, for a simple graph G , to be the size of the largest “homogeneous subset” of G , i.e., a clique or an independent set. One way to state Ramsey’s Theorem is to say that $\text{hom}(G) \gg \log(n)$ for G on n vertices.

Conjecture 1 (Erdős, Hajnal [5]). For every graph H , there is an $\epsilon > 0$ so that, if G on n vertices has no induced copy of H , then $\text{hom}(G) \gg n^\epsilon$.

The intuition behind this conjecture is that the property of avoiding H as an induced subgraph is a mark of being very far from random, since a random graph will have many copies of H . On the other hand, it is notoriously difficult to deterministically construct a graph G coming anywhere close to the Ramsey bound. In a sense, $\text{hom}(G)$ is a statistic which is *very* sensitive to “true” randomness. Therefore, if G avoids H , then $\text{hom}(G)$ should be much larger than $\log(n)$ – something like n^ϵ for $\epsilon > 0$.

The Erdős-Hajnal conjecture, as stated above, does not directly apply to our situation, however, since we are interested in tournaments instead of graphs. Fortunately, Alon, Pach, and Solymosi ([2]) have provided a very clever argument that the following analogous statement is actually equivalent to Conjecture 1. Define $\text{trans}(T)$ to be the size of the largest transitive subtournament of the tournament T .

Conjecture 2. For every tournament T , there is an $\epsilon > 0$ so that, if T' on n vertices has no copy of T , then $\text{trans}(T') \gg n^\epsilon$.

Therefore, if we believe Conjecture 1, then the fact that a tournament arising from a line configuration cannot contain a copy of T_0 implies that there is some $\epsilon_{\text{LINE}} > 0$ so that every configuration of lines in \mathbb{R}^3 contains a “stacked” subset of size $n^{\epsilon_{\text{LINE}}}$. Indeed, in the same way, the Erdős-Hajnal Conjecture coupled with the Milnor-Thom Theorem implies that *any* tournament on a set of points in \mathbb{R}^d whose edges are determined by the sign of a polynomial in the coordinates of the endpoints contains such a “polynomial-sized” transitive subtournament.

Recently, a even more general conjecture by Babai in [3] was resolved unconditionally by Alon, Pach, Pinchasi, Radoičić, and Sharir ([1]). Their powerful result, which has endless applications, is as follows:

Theorem 2. Let \mathcal{F} be a family of n semi-algebraic sets in \mathbb{R}^d of constant description complexity, and let $R \subset \mathcal{F} \times \mathcal{F}$ be a fixed semi-algebraic relation on \mathcal{F} . Then there exists a constant $\epsilon > 0$, which depends only on the maximum description complexity of the sets in \mathcal{F} and of R , and a subfamily $\mathcal{F}' \subset \mathcal{F}$ with at least n^ϵ elements, such that either every pair of distinct elements of \mathcal{F}' belongs to R , or no such pair belongs to R .

In the case of line configuration tournaments, they are able to show that $\epsilon_{\text{LINE}} \geq 1/6$. In the other direction, we have:

Proposition 3. $\epsilon_{\text{LINE}} \leq \log 3 / \log 7 \approx 0.564575$.

The proof of this fact relies on a computation and the following observation. Consider the three “non-stacked” lines of Figure 1. By applying a linear transformation, we may “bundle” these lines into the union of a (double) cone and a cylinder, as thin as we wish, without disturbing the crossing relations. Then, the three cones can be arranged in the same way as the lines of the original three lines, so that now we have 9 lines with no stacked subset of size larger than 4. (See Figure 2.) Proceeding recursively, we can construct a family of 3^k lines with no stacked subset of size larger than 2^k , i.e., $\epsilon_{\text{LINE}} \leq \log 2 / \log 3$.




Figure 2: Bundling lines together recursively.

An extensive computer search revealed the existence of a configuration of 7 lines with largest stacked subset of size 3. Therefore, we can improve this bound by the same construction to $\epsilon_{\text{LINE}} \leq \log 3 / \log 7$. \square

Given a real-valued function f on \mathbb{R}^d and a set of points $V \subset \mathbb{R}^d$, define the digraph $\Gamma^f(V)$ to have V as its vertex set and $(x, y) \in E(\Gamma^f(V))$ iff $f(y - x) > 0$. If f is odd, then $\Gamma^f(S)$ is a tournament; if f is even, we may consider it a simple graph. By “complete graph”, we mean the digraph whose edges are all pairs (u, v) with u and v any two distinct vertices. We call a set $V \subset \mathbb{R}^d$ “ f -general” if $f(v_1 - v_2) \neq 0$ for any $v_1, v_2 \in V$. Call a function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ “Erdős-Hajnal” if there exists an $\epsilon > 0$ so that, for any f -general $V \subset \mathbb{R}^d$ with $|V| = n$, $\Gamma^f(V)$ contains a complete graph, an independent set, or an induced transitive subtournament of cardinality $\gg n^\epsilon$. The requirement that V be f -general corresponds to the requirement that the lines in a

configuration be skew. This assumption, though not strictly necessary, simplifies the discussion below.

It is also possible to generalize this idea as follows. For a set $S \subset \mathbb{R}^d$ and a set of points $V \subset \mathbb{R}^d$, define $\Gamma^S(V)$ to have V as its vertex set and $(x, y) \in E(\Gamma^S(V))$ iff $y - x \in S$. Clearly, the definition of $\Gamma^f(V)$ coincides with this definition when $S = f^{-1}(\mathbb{R}^+)$, at least for f -general sets V . Furthermore, we say that S is “Erdős-Hajnal” if there exists an $\epsilon > 0$ so that, for any $V \subset \mathbb{R}^d$ with $|V| = n$, $\Gamma^S(V)$ contains either a complete graph, an independent set, or an induced transitive subtournament of cardinality $\gg n^\epsilon$.

Our main question here is: which functions/sets are Erdős-Hajnal? By Theorem 2, all polynomials f and any semi-algebraic set S are Erdős-Hajnal. In the next section, we prove some basic facts and describe a much larger class of Erdős-Hajnal sets. In particular, we show that all sets belonging to the set algebra \mathfrak{A}_d generated by semigroups in \mathbb{R}^d are Erdős-Hajnal. Therefore, in light of Theorem 2, it is a natural question to ask whether semi-algebraic sets belong to \mathfrak{A}_n . In Section 2, we prove that this is the case for $k \leq 2$ and discuss the situation for $k \geq 3$.

1 Erdős-Hajnal Sets

We collect some basic results concerning Erdős-Hajnal sets. First, the set algebra generated by Erdős-Hajnal sets consists entirely of Erdős-Hajnal sets. For $X \subset \mathbb{R}^d$, write \bar{X} for the topological closure of X and X^c for the complement of X .

Proposition 4. Complement, intersection, and union preserve the property of being Erdős-Hajnal.

Proof. The first claim is obvious. For the second, let A and B be Erdős-Hajnal sets, and consider $G = \Gamma^{A \cap B}(V)$ for some set V in \mathbb{R}^d of cardinality n . The vertex set of $G_1 = \Gamma^A(V)$ contains a subset V_0 of size n^{ϵ_A} which induces an independent set, a complete graph, or a transitive tournament. Since B is Erdős-Hajnal, $\Gamma^B(V_0)$ has a subset V_1 of size $n^{\epsilon_{A \cap B}}$ which induces an empty, complete, or transitive digraph G_2 . Since $G' = \Gamma^{A \cap B}(V_1) = \Gamma^A(V_1) \cap \Gamma^B(V_1)$, G' is empty if either G_1 or G_2 is. If either G_i is complete, then G' is empty, complete, or transitive according to whether the other G_j is. If both G_1 and G_2 are transitive tournaments, then, by the Erdős-Szekeres Theorem, their intersection contains either a transitive subtournament of size $n^{\epsilon_{A \cap B}/2}$ or an independent set of size $n^{\epsilon_{A \cap B}/2}$.

The fact that the union of two Erdős-Hajnal sets is Erdős-Hajnal now follows from De Morgan’s Law. \square

Corollary 5. If f and g are Erdős-Hajnal functions, then $f \cdot g$ is also.

Proof. The semialgebraic set $(fg)^{-1}(\mathbb{R}^+)$ is the complement of the symmetric difference of the sets $A = f^{-1}(\mathbb{R}^+)$ and $B = g^{-1}(\mathbb{R}^+)$, i.e., $(A \cap B) \cup (A^c \cap B^c)$, except for some points of $f^{-1}(0)$ and $g^{-1}(0)$. These sets are of no consequence, however, since fg -generality implies f -generality and g -generality. \square

Our next results says that any *bounded* set which does not have the origin on its boundary is Erdős-Hajnal.

Proposition 6. Any bounded set $B \subset \mathbb{R}^d$ with $\mathbf{0} \notin \partial B$ is Erdős-Hajnal.

Proof. First, suppose that $\mathbf{0} \in \bar{B}$. Since $\mathbf{0} \notin \partial B = \bar{B} \setminus \text{int } B$, it must be that $\mathbf{0} \in \text{int } B$. Therefore, there is some $r > 0$ so that the closed ball of radius r about the origin is contained in B , and an $R \geq r$ so that the closed ball of radius R about the origin contains B . Partition space into a lattice of axis-parallel hypercubes (i.e., $[0, s]^d$) with side length $s > 0$ so that $s\sqrt{d} \leq r$. Let Λ denote the set of cubes containing the points of the lattice generated by $t\mathbf{e}_1, \dots, t\mathbf{e}_d$, where $t = s(\lceil R/s \rceil + 1)$. The t^d distinct “cosets” $\Lambda_1, \dots, \Lambda_{t^d}$ of Λ partition the points of V , and no such Λ_i contains a pair x, y with $r < |y - x| \leq R$. Some Λ_i contains at least $n' = n/t^d$ points of V , and, for each $x, y \in \Lambda_i$, either there are $\sqrt{n'}$ points $V_0 \subset V$ in one of the cubes comprising Λ_i , or there are $\sqrt{n'}$ points $V_0 \subset V$ in distinct cubes of Λ_i . In the former case, $y - x \in B$ for every pair of points in V_0 , so V_0 induces a complete digraph. In the latter case, $y - x \notin B$ for every pair $x, y \in V_0$, so V_0 induces an independent set.

On the other hand, if $\mathbf{0} \notin \bar{B}$, then $\mathbf{0} \in \text{int}(B^c)$. There are numbers r and R , $0 < r < R$, so that the closed ball U of radius R about the origin contains B and the closed ball U' of radius r about the origin is disjoint from B . Let $A = U \setminus B$, so $U' \subset A$. A and U are bounded sets whose closure contains $\mathbf{0}$, but whose boundary does not. We may therefore apply the first part of the proof to each of these sets, and invoke Proposition 4 by noting that $B = U \setminus A$. \square

The last result of this section is that any *semigroup* – a subset of \mathbb{R}^d which is closed under addition – is Erdős-Hajnal.

Proposition 7. Any semigroup S is Erdős-Hajnal.

Proof. By Proposition 4, we may assume that S lies in one orthant. In that case, note that $S \cap -S = \emptyset$. Furthermore, if $(x, y) \in \Gamma^S(V)$ and $(y, z) \in \Gamma^S(V)$, then $y - x \in S$ and $z - y \in S$, so $z - x = (z - y) + (y - x) \in S$. Therefore, if we define $y \succ x$ for $x \neq y$ whenever $(x, y) \in \Gamma^S(V)$, then \succ is antisymmetric and transitive, i.e., a partial order. By Dilworth’s Theorem, there is a subset of V of size $n^{1/2}$ which is either a chain or an antichain, i.e., induces a transitive subtournament of $\Gamma^S(V)$ or is an independent set. \square

2 Dimension at Most Two

Since every element of the set algebra \mathfrak{A}_d generated by semigroups in \mathbb{R}^d is Erdős-Hajnal, it is natural to ask whether all semi-algebraic sets belong to this family. If so, this would give another proof of Theorem 2. Moreover, we consider this question interesting in its own right. In this section, we prove the following Theorem.

Theorem 8. Every semi-algebraic set in dimension $d \leq 2$ belongs to \mathfrak{A}_d .

Note that the statement is obvious in the case that f is a function of only one variable: S is a finite union of intervals whose (possibly infinite) endpoints are on the same side of the origin, and each such interval is either a semigroup itself or a difference of two semigroups. Also, by Corollary 5, it suffices to prove the theorem for sets defined by irreducible polynomials. We begin with the following Lemma.

Lemma 9. Any irreducible polynomial f in two variables for which there are infinitely many solutions to $\langle(x, y), \nabla f(x, y)\rangle = 0$ with $f(x, y) = 0$ must be of the form $f(x, y) = ax + by$ for some a and b .

Proof. Suppose that there are infinitely many pairs (x, y) with $\langle(x, y), \nabla f(x, y)\rangle = 0$ and $f(x, y) = 0$. By Bézout's Theorem, the function

$$h(x, y) = \langle(x, y), \nabla f(x, y)\rangle = x \frac{\partial f}{\partial x} + y \frac{\partial f}{\partial y}$$

must be divisible by $f(x, y)$. Consider the terms of highest x -degree in f , and choose among them the term of highest y degree, say $cx^r y^s$. It is easy to see that the analogous term for h is $(r+s)cx^r y^s$. Since these terms are unique, $(r+s)c = c$, i.e., $r+s = 1$. Therefore, $f = ax + g(y)$ for some single-variable polynomial g and constant a . By symmetry, then, f must be of the form $f = ax + by + d$. However, f cannot have any constant term, since h has no constant term, so $d = 0$. \square

Lemma 10. Any irreducible polynomial f in two variables for which there are infinitely many solutions to $\langle(-y, x), \nabla f(x, y)\rangle = 0$ with $f(x, y) = 0$ must be of the form $f(x, y) = a(x^2 + y^2) + b$ for some a and b .

Proof. Suppose that there are infinitely many common solutions to $f(x, y) = 0$ and $\langle(-y, x), \nabla f(x, y)\rangle = 0$. By Bézout's Theorem, the function

$$h(x, y) = \langle(-y, x), \nabla f(x, y)\rangle = x \frac{\partial f}{\partial y} - y \frac{\partial f}{\partial x}$$

must be divisible by $f(x, y)$. Write $h = g \cdot f$. The total degree of h is the same as the total degree of f , so $g(x, y) = b$, a constant. Furthermore, the operator $x D_y - y D_x$ preserves total degree. Fix a degree d , denote by c_i the coefficient of $x^i y^{d-i}$ in f , and denote by c'_i the coefficient of $x^i y^{d-i}$ in h . Then

$$\begin{bmatrix} c'_0 \\ c'_1 \\ \vdots \\ c'_{d-1} \\ c'_d \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 & 0 & 0 \\ -d & 0 & 2 & \cdots & 0 & 0 & 0 \\ 0 & 1-d & 0 & \ddots & 0 & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \ddots & 0 & d-1 & 0 \\ 0 & 0 & 0 & \cdots & -2 & 0 & d \\ 0 & 0 & 0 & \cdots & 0 & -1 & 0 \end{bmatrix} \begin{bmatrix} c_0 \\ c_1 \\ \vdots \\ c_{d-1} \\ c_d \end{bmatrix} = b \cdot \begin{bmatrix} c_0 \\ c_1 \\ \vdots \\ c_{d-1} \\ c_d \end{bmatrix}.$$

Therefore, b is an eigenvalue of this matrix, which we will denote M_d . We argue that the (nonzero) eigenvalues of M_d are pure imaginary. Let D be the $d \times d$ diagonal

matrix whose $(k+1)^{\text{th}}$ diagonal entry is $i^k \sqrt{\binom{d}{k}}$. Then $M'_d = D^{-1}M_dD$ is a symmetric tridiagonal matrix whose diagonal entries are all zero and whose k^{th} super- or subdiagonal element is $ik(-1)^{k+1} \sqrt{\binom{n}{k-1}} \binom{n}{k}$. Since M'_d is symmetric pure imaginary, its eigenvalues – and therefore the eigenvalues of M_d – are pure imaginary. This contradicts the fact that the c'_j and c_j are all real unless $b = 0$.

Now, the last d columns of M_d are linearly independent, so the nullspace of M_d has dimension at most one. On the other hand, the negative of each eigenvalue $\lambda \in \sigma(M_d)$ is also an eigenvalue, since

$$M_d(J\mathbf{v}) = (M_dJ)\mathbf{v} = -(JM_d)\mathbf{v} = -J(M_d\mathbf{v}) = -J\lambda\mathbf{v} = -\lambda(J\mathbf{v}),$$

where J is the “exchange” matrix whose only nonzero entries are ones on the antidiagonal, and \mathbf{v} is an eigenvector corresponding to λ . Therefore, the number of nonzero elements of $\sigma(M_d)$ is even. We may conclude that M_d is invertible if d is odd, and has nullity 1 if d is even. In the latter case, it is easy to verify that the $(d+1)$ -vector \mathbf{w} whose k^{th} element is 0 if k is even and $\binom{d/2}{(k-1)/2}$ if k is odd is a null vector of M_d . The polynomial corresponding to \mathbf{w} is $(x^2 + y^2)^{d/2}$, so we may write $f = f_0(x^2 + y^2)$ for some polynomial f_0 .

If the factorization of f_0 over \mathbb{R} has no linear terms, then $f^{-1}(\mathbb{R}^+) = \emptyset$, a contradiction. Therefore, $f_0(z)$ is divisible by $z + a$ for some $a \in \mathbb{R}$, and f is divisible by $x^2 + y^2 + a$. Since f is irreducible, the lemma follows. \square

Lemma 11. For any irreducible polynomial $f(x, y)$ in two variables which is not of the form $ax + by$ or $a(x^2 + y^2) + b$ for any a, b , there is a family $\{I_j\}_{j=1}^m$ of open intervals of $[0, 2\pi]$, with $|I_j| \leq \pi/2$ for each j , and a corresponding family $\{g_j\}_{j=1}^m$ of C^∞ , monotone functions defined on each of the I_j 's respectively, so that the variety defined by f – i.e., $\{(x, y) : f(x, y) = 0\}$ – is the disjoint union of the graphs of the g_j in polar coordinates, with the exception of finitely many points.

Proof. The set X of zeroes of f is a one-dimensional affine variety with finitely many singularities and finitely many components (in the Euclidean topology). (See [4] for basic facts concerning the differential topology of real algebraic varieties.) Furthermore, by the implicit function theorem, removing the singularities and $\mathbf{0}$ from X leaves a finite number of C^∞ arcs, i.e., diffeomorphic images of open intervals. (We consider a topological S^1 an arc in the obvious way, i.e., by removing one point.) Since f is not divisible by x or y , X has finitely many points in common with the x and y -axes, so we may even remove the two axes and still obtain such a decomposition. Clearly, it suffices to prove the Lemma for a single one of the resulting arcs, say, $\{(x(t), y(t)) : t \in (0, 1)\}$. We may also assume, without loss of generality, that all the points of this arc lie in the first quadrant.

Consider the function $\phi(t) = \tan^{-1}(y(t)/x(t))$. Differentiating yields

$$\phi'(t) = \frac{y'x - x'y}{y^2 + x^2}.$$

Suppose there are infinitely many $t \in (0, 1)$ for which $\phi'(t) = 0$. Then $y'x - x'y = 0$ at each such t , and multiplying by $\partial f / \partial x$ and applying the identity $x' \cdot \partial f / \partial x + y' \cdot$

$\partial f / \partial y = 0$ (obtained by differentiating $f(x, y) = 0$),

$$\begin{aligned} 0 &= \frac{\partial f}{\partial x}(y'x - x'y) = \frac{\partial f}{\partial x}y'x - \frac{\partial f}{\partial x}x'y \\ &= \frac{\partial f}{\partial x}y'x + \frac{\partial f}{\partial y}y'y \\ &= y' \cdot \langle (x, y), \nabla f(x, y) \rangle. \end{aligned}$$

By symmetry, $x' \langle (x, y), \nabla f(x, y) \rangle = 0$. However, x' and y' cannot both be zero, so $\langle (x, y), \nabla f(x, y) \rangle = 0$. Lemma 9 implies that $f(x, y) = ax + by$ for some a and b , contradicting our hypothesis. Hence, there exist only a finite set $\{t_j\}_{j=1}^l$ of values of $t \in (0, 1)$ for which $\phi'(t) = 0$.

Now, consider the quantity $d(x^2 + y^2)/d\phi = 2(xx' + yy')(x^2 + y^2)/(y'x - x'y)$. If this equals zero at $t \notin \{t_j\}_{j=1}^l$, then $xx' + yy' = 0$. Again, applying the identity $x' \cdot \partial f / \partial x + y' \cdot \partial f / \partial y = 0$ and noting that x' and y' cannot simultaneously be zero, we obtain that

$$x \frac{\partial f}{\partial y} - y \frac{\partial f}{\partial x} = 0.$$

By Lemma 10, there are only finitely many t'_j for which this equality can hold.

If we remove the points $(x(t_j), y(t_j))$ and $(x(t'_j), y(t'_j))$ from the arc, we are left with a finite collection of arcs on which ϕ is C^∞ , monotone, and defined on an open interval of length at most $\pi/2$. Choosing one such interval I and letting

$$g(\theta) = \sqrt{(x \circ (\phi|_I)^{-1}(\theta))^2 + (y \circ (\phi|_I)^{-1}(\theta))^2}$$

gives one of the g_j of the desired decomposition: g_j is monotone because

$$\frac{dg_j^2}{d\theta} = 2 \left((x \circ (\phi|_I)^{-1}(\theta)) \cdot \frac{dx}{d\phi} \Big|_{\phi=\theta} + (y \circ (\phi|_I)^{-1}(\theta)) \cdot \frac{dy}{d\phi} \Big|_{\phi=\theta} \right) \cdot \frac{d\phi^{-1}}{d\theta} \neq 0.$$

The other g_j are obtained analogously. \square

Proof of Theorem 8. First, if $f(x, y) = ax + by$, then S is a half-plane, which is a semigroup, and the result follows immediately from Proposition 7. If $f(x, y) = a(x^2 + y^2) + b$, then either S is empty (in which case the result is immediate) or else S^G is a disk (in which case S is the union of four semigroups). Otherwise, let $\{g_j\}_{j=1}^N$ be the collection of all the g_j constructed in the previous Lemma, and let $\{I_j\}_{j=1}^N$ be the corresponding intervals. Some of the g_j we will classify as “unimportant” as follows. For each point \mathbf{x}_t on the graph Γ_j of g_j – i.e., the image of $t \in (0, 1)$ in polar coordinates – consider the quantities

$$T_-(t) = \lim_{\lambda \rightarrow 1^-} \operatorname{sgn}(f(\lambda \mathbf{x}_t)) \text{ and } T_+(t) = \lim_{\lambda \rightarrow 1^+} \operatorname{sgn}(f(\lambda \mathbf{x}_t)).$$

We argue these limits exist and equal ± 1 . If they did not exist (or equalled 0), then, since sgn is continuous everywhere but at 0, there must be an infinite sequence of $\lambda_i \rightarrow 1-$ (say) so that $f(\lambda_i \mathbf{x}_t) = 0$. But then, by Bézout’s Theorem, f must be

divisible by the equation of the line passing through \mathbf{x}_t and the origin, a contradiction. Furthermore, T_- and T_+ are continuous functions of t : suppose, to the contrary, that there is a point $\mathbf{x}_t \in \Gamma_j$ and two sequences t_i^1 and t_i^2 so that $T_-(t_i^1) = 1$ and $T_-(t_i^2) = -1$, $t_i^1, t_i^2 \rightarrow t$. Then there are sequences λ_i^1 and a λ_i^2 so that $\lambda_i^1, \lambda_i^2 \rightarrow 1-$, $\text{sgn}(f(\lambda_i^1 x_i^1)) = +1$ and $\text{sgn}(f(\lambda_i^2 x_i^2)) = -1$. For each i , we interpolate between $\lambda_i^1 x_i^1$ and $\lambda_i^2 x_i^2$ with the function $h_i : [t_i^1, t_i^2] \rightarrow \mathbb{R}^2$ defined by $h_i(s) = \lambda_i(s)\mathbf{x}_s$ where

$$\lambda_i(s) = \frac{\lambda_i^2 - \lambda_i^1}{t_i^2 - t_i^1}(s - t_i^1) + \lambda_i^1.$$

Since $f(h_i(t_i^1)) > 0$ and $f(h_i(t_i^2)) < 0$, there is an s_i so that $f(h_i(s_i)) = 0$, by the intermediate value theorem. Furthermore, it is easy to see that $s_i \rightarrow t$ and $\lambda_i(s_i) \rightarrow 1-$. Then there is a sequence of points of $V = f^{-1}(0)$ not lying on the curve g_j , but approaching $g_j(t)$, implying that $g_j(t)$ is a singular point of V (a self-intersection, in particular). This contradicts the fact that g_j contains no singularities of V . Therefore, we may conclude that T_- and T_+ are continuous. Since they take on the values ± 1 , however, this means that they are constant. Hence, we may write c_j^- for the value of T_- on g_j and c_j^+ analogously for T_+ . Finally, we say that g_j is “unimportant” if $c_j^- = c_j^+$, and re-index the g_j so that all the elements of $\{g_j\}_{j=1}^M$ are “important”.

Now, let X_j denote the set $\{\lambda g_j(\theta) \text{ cis}(\theta) : \lambda \geq 1, \theta \in I_j\}$ if $c_j^- = +1$ and $\{\lambda g_j(\theta) \text{ cis}(\theta) : \lambda > 1, \theta \in I_j\}$ if $c_j^- = -1$, where $\text{cis}(\theta) = (\cos(\theta), \sin(\theta))$; let Y be the set of $\mathbf{v} \in \mathbb{R}^2 \setminus \{\mathbf{0}\}$ so that for all sufficiently small $\epsilon > 0$ we have $f(\epsilon\mathbf{v}) > 0$; let Y' be the set of $\mathbf{v} \in \mathbb{R}^2 \setminus \{\mathbf{0}\}$ so that for all sufficiently small $\epsilon > 0$ we have $f(\epsilon\mathbf{v}) < 0$; and define Z to be $Y \cdot \mathbb{R}^+$. Define A to be the set of points lying on the graph of some g_j , let B be the (finite) set of points $f^{-1}(0) \setminus A$, let $C = B \cdot \mathbb{R}^+$, and let $D = f^{-1}(\mathbb{R}^+) \cap C$. We argue that

$$S' = S \setminus \{\mathbf{0}\} = ((Z \ominus X_1 \ominus \cdots \ominus X_M) \setminus C) \cup D, \quad (1)$$

where “ \ominus ” denotes symmetric difference (which is associative). Denote the set $X_1 \ominus \cdots \ominus X_M$ by W , and consider a ray $\rho_{\mathbf{v}} = \mathbf{v} \cdot \mathbb{R}^+$ with $\mathbf{v} \notin C$. Let $h_{\mathbf{v}}(\lambda) = f(\lambda\mathbf{v})$ for $\lambda \geq 0$. Since f is continuous and the graphs of the g_j completely cover the set $f^{-1}(0) \cap C$, $h_{\mathbf{v}}$ undergoes a sign change at λ iff $\lambda\mathbf{v}$ lies on the graph of some “important” g_j . Therefore, $S' \cap \rho_{\mathbf{v}} = \rho_{\mathbf{v}} \cap W^C$ whenever $\mathbf{v} \in Y$. On the other hand, if $\mathbf{v} \notin Y$, then $\mathbf{v} \in Y'$. If not, then f has zeroes arbitrarily close to the origin along the ray $\rho_{\mathbf{v}}$, which we previously argued implies that $f(x, y) = ax + by$ for some a and b . It is clear then that $S' \cap \rho_{\mathbf{v}} = W \cap \rho_{\mathbf{v}}$ if $\mathbf{v} \notin Y$, so that $S' \setminus C = (Z \ominus W) \setminus C$.

Define $\eta(\theta)$ to be 1 if $\text{cis}(\theta) \in Y$ and -1 if $\text{cis}(\theta) \in Y'$, and let $l(\theta) = \min\{1, \epsilon\}$, where ϵ is the supremum of the set of reals so that $0 < \epsilon' < \epsilon$ implies

$$(\text{sgn} \circ f)(\epsilon' \cdot \text{cis}(\theta)) = \eta(\theta).$$

Since all the ϵ are positive and the unit circle is compact, there is a δ so that $l(\theta) > \delta$ for all $\theta \in [0, 2\pi]$. Therefore, $\eta(\theta) = (\text{sgn} \circ f)(\delta \cdot \text{cis}(\theta))$ for all θ . If η has infinitely many discontinuities, then f has infinitely many zeroes on the circle of radius δ about

the origin, which in turn implies (as previously argued) that $f(x, y) = a(x^2 + y^2) + b$ for some a and b . Hence, $\eta^{-1}(1)$ is a finite collection of intervals, and

$$Z = \{\lambda \cdot \text{cis}(\theta) : \lambda > 0, \theta \in \eta^{-1}(1)\}$$

is a finite union of cones (projective hulls of convex sets), each of which is a semigroup.

Now, suppose $\mathbf{x}, \mathbf{y} \in X_j$, so that $\mathbf{x} = \lambda_1 g_j(\theta_1) \text{cis}(\theta)$ and $\mathbf{y} = \lambda_2 g_j(\theta_2) \text{cis}(\theta)$ with $|\theta_1 - \theta_2| \leq \pi/2$, $\theta_1, \theta_2 \in [0, 2\pi]$, and $\lambda_1, \lambda_2 \geq 1$. We may assume without loss of generality that \mathbf{x} and \mathbf{y} both lie in the first quadrant. If we write $\mathbf{x} + \mathbf{y}$ as (r, ϕ) in polar coordinates, then it is easy to see that $\theta_1 < \phi < \theta_2$ and $r \geq \max\{\lambda_1 g_j(\theta_1), \lambda_2 g_j(\theta_2)\}$. By the monotonicity of g_j , $r > g_j(\phi)$. If $\mathbf{x} + \mathbf{y} \notin X_j$, then $r \leq g_j(\phi)$, a contradiction. Hence, X_j is a semigroup.

That C is a finite union of semigroups is obvious. We can write D as

$$D = \bigcup_{\mathbf{v} \in B} (f_0^{-1}(\mathbb{R}^+) \cap \rho_{\mathbf{v}}).$$

Because $f_0(\lambda) = f(\lambda \mathbf{v})$ is a polynomial in a single variable, f_0 is Erdős-Hajnal. Furthermore, $(f_0^{-1}(\mathbb{R}^+) \cdot \mathbf{v}) \cap \rho_{\mathbf{v}} = f^{-1}(\mathbb{R}^+) \cap \rho_{\mathbf{v}}$, and $\rho_{\mathbf{v}}$ is a semigroup, so $D \in \mathfrak{A}_2$ since $f_0^{-1}(\mathbb{R}^+) \in \mathfrak{A}_1$ implies $f_0^{-1}(\mathbb{R}^+) \cdot \mathbf{v} \in \mathfrak{A}_2$.

Finally, since each of the terms in (1) belongs to \mathfrak{A}_2 , we may apply Proposition 4 to conclude that S is Erdős-Hajnal. \square

One is tempted to try to generalize Theorem 8 to higher dimensions. However, we believe this not to be possible, at least for dimension greater than three. We have the following (perhaps premature) conjecture.

Conjecture 3. There exist semi-algebraic sets in dimension $d \geq 4$ which do not belong to \mathfrak{A}_d , but all semi-algebraic sets defined by three-variable polynomials *do* belong to \mathfrak{A}_3 .

In fact, we believe the function $f(a, b, c, d) = ad - bc$ provides a counterexample for dimension four. Ironically, showing that f is Erdős-Hajnal is precisely what is needed to show that $\epsilon_{\text{LINE}} > 0$.

3 Acknowledgements

Thank you to János Pach, Ricky Pollack, and Uli Wagner for helpful discussions and ideas. And a special thank you to József Solymosi for his insight and guidance, as well as for introducing me to this wonderful area of research.

References

- [1] N. Alon, J. Pach, R. Pinchasi, R. Radoičić, M. Sharir, Crossing patterns of semi-algebraic sets. *J. Comb. Theory Ser. A* (2004), to appear.

- [2] N. Alon, J. Pach, J. Solymosi, Ramsey-type theorems with forbidden subgraphs. *Combinatorica* 21 (2001), no. 2, 155–170.
- [3] L. Babai, Open problem, in: Proc. of the Conference in Keszthely, Hungary, 1976, vol. 2, North Holland (1978), 1189.
- [4] J. Bochnak, M. Coste, M.-F. Roy, Real algebraic geometry. *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*, 36. Springer-Verlag, Berlin, 1998.
- [5] P. Erdős, A. Hajnal, Ramsey-type theorems. *Discrete Appl. Math.* 25 (1989), no. 1-2, 37–52.
- [6] P. Erdős, A. Hajnal, J. Pach, A Ramsey-type theorem for bipartite graphs. *Geombinatorics* 10 (2000), no. 2, 64–68.
- [7] J. Spencer, The strange logic of random graphs. *Algorithms and Combinatorics*, 22. Springer-Verlag, Berlin, 2001.
- [8] H. E. Warren, Lower bounds for approximation by nonlinear manifolds. *Trans. Amer. Math. Soc.* 133 (1968), 167–178.