

Counting rooted spanning forests in complete multipartite graphs

László A. Székely¹
 Department of Mathematics
 University of South Carolina
 Columbia, SC 29208

Abstract

Jin and Liu discovered an elegant formula for the number of rooted spanning forests in the complete bipartite graph K_{a_1, a_2} , with b_1 roots in the first vertex class and b_2 roots in the second vertex class. We give a simple proof to their formula, and a generalization for complete m -partite graphs, using the multivariate Lagrange inverse.

Y. Jin and C. Liu [3] give a formula for $f(m, l; n, k)$, the number of spanning forests of the labelled complete bipartite graph $K_{n, m}$, where in the forest every tree is rooted, there are k roots in the first vertex class (among the n vertices) and l roots in the second vertex class (among the m vertices), and the trees in the forest are not ordered. They discovered the elegant formula

$$f(m, l; n, k) = \binom{m}{l} \binom{n}{k} n^{m-l-1} m^{n-k-1} (km + ln - lk). \quad (1)$$

The goal of the present note is generalization of (1) from complete bipartite to complete multipartite graphs, through a simple proof using the multivariate Lagrange inverse.

Let $f(a_1, b_1; \dots; a_m, b_m)$ denote the number of spanning forests of the labelled complete multipartite graph K_{a_1, a_2, \dots, a_m} , where in the forest every tree is rooted, there are b_i roots in the i^{th} vertex class for $i = 1, 2, \dots, m$, and the trees in the forest are not ordered. Let $w_i(t_1, \dots, t_m)$ denote the multivariate exponential generating function (EGF) of the numbers $f(a_1, 0; \dots; a_i, 1; \dots; a_m, 0)$ (the number of rooted spanning trees of the complete multipartite graph K_{a_1, a_2, \dots, a_m} , if the root has to be in the i^{th} class), i.e.

$$w_i(t_1, \dots, t_m) = \sum_{a_1=0}^{\infty} \dots \sum_{a_i=1}^{\infty} \dots \sum_{a_m=0}^{\infty} f(a_1, 0; \dots; a_i, 1; \dots; a_m, 0) \prod_{k=1}^m \frac{t_k^{a_k}}{a_k!}. \quad (2)$$

¹Research partially supported by the NSF Grant 0072187 and the Hungarian NSF Grant T 032455.

The key identity for our argument is

$$t_i e^{(w_1+w_2+\dots+w_m)-w_i} = w_i \quad \text{for } i = 1, 2, \dots, m. \quad (3)$$

The proof of formula (3) is based on the following combinatorial decomposition. Given a rooted spanning tree of the complete multipartite graph K_{a_1, a_2, \dots, a_m} , where the root is in the i^{th} class, remove the root vertex from the tree to obtain a spanning forest of $K_{a_1, a_2, \dots, a_i-1, \dots, a_m}$, and mark the former neighbors of the eliminated root vertex as roots in the forest. This decomposition establishes a bijection between the following two sets:

the set of rooted spanning trees of the complete multipartite graph K_{a_1, a_2, \dots, a_m} , where the root is in the i^{th} vertex class,

and

the set of some ordered pairs, where the first entry of the ordered pair is one of the vertices of the i^{th} vertex class, the second element of the ordered pair is a rooted spanning forest of $K_{a_1, a_2, \dots, a_i-1, \dots, a_m}$, where the vertex from the first entry is removed from the i^{th} vertex class, and the trees of the forest are not ordered.

Now $t_i e^{(w_1+w_2+\dots+w_m)-w_i}$ is the EGF of the set of ordered pairs in question, according to the Exponential Formula; and w_i is the same EGF by the bijection. Set $\Phi_i(w_1, w_2, \dots, w_m) = e^{(w_1+w_2+\dots+w_m)-w_i}$.

According to the multiplication rule of EGF's, $\prod_{k=1}^m w_k^{b_k}$ is the multivariate exponential generating function of the number of rooted spanning forests of complete m -partite graphs, with b_k roots in the k^{th} vertex class, where the trees rooted in the same part *are ordered*; hence

$$f(a_1, b_1; \dots; a_m, b_m) = \frac{a_1! a_2! \cdots a_m!}{b_1! b_2! \cdots b_m!} [t_1^{a_1} t_2^{a_2} \cdots t_m^{a_m}] \prod_{k=1}^m w_k^{b_k}. \quad (4)$$

According to Part 1 of Theorem 1.2.9 (Multivariate Lagrange Formula) from [2], (3) implies

$$[t_1^{a_1} t_2^{a_2} \cdots t_m^{a_m}] \prod_{k=1}^m w_k^{b_k} = [\lambda_1^{a_1} \cdots \lambda_m^{a_m}] \left\{ \det \left| \delta_{ij} - \frac{\lambda_j}{\Phi_i} \cdot \frac{\partial \Phi_i}{\partial \lambda_j} \right| \right. \quad (5)$$

$$\left. \times \lambda_1^{b_1} \cdots \lambda_m^{b_m} \prod_{k=1}^m e^{a_i(w_1+\dots+w_m)-a_i w_i} \right\}, \quad (6)$$

where Φ_i is a short-hand notation for $\Phi_i(\lambda_1, \dots, \lambda_m)$. Observe that $\frac{\lambda_j}{\Phi_i} \cdot \frac{\partial \Phi_i}{\partial \lambda_j} = (1 - \delta_{ij})\lambda_j$, and the for the determinant in (5) we have the well-known evaluation

$$\det \left| \delta_{ij} - (1 - \delta_{ij})\lambda_j \right| = (\lambda_1 + 1) \cdots (\lambda_m + 1) \left(1 - \frac{\lambda_1}{\lambda_1 + 1} - \cdots - \frac{\lambda_m}{\lambda_m + 1} \right) \quad (7)$$

(see for example Exercise 225 in [1]). Now (7) is easily rewritten as

$$1 - \sum_{j=2}^m (j-1) \sum_{1 \leq i_1 < i_2 < \cdots < i_j \leq m} \lambda_{i_1} \lambda_{i_2} \cdots \lambda_{i_j}, \quad (8)$$

and (8) is rewritten as

$$\sum_{l_1=0}^1 \sum_{l_2=0}^1 \dots \sum_{l_m=0}^1 (1 - l_1 - l_2 - \dots - l_m) (\lambda_1)_{l_1} (\lambda_2)_{l_2} \dots (\lambda_m)_{l_m}, \quad (9)$$

where $(x)_t$ stands for the falling factorial, $(x)_0 = 1$ and $(x)_1 = x$. Introducing the notation $A = a_1 + a_2 + \dots + a_m$ and using (9), we find that (5) and (6) are equal to

$$\begin{aligned} & [\lambda_1^{a_1-b_1} \lambda_2^{a_2-b_2} \dots \lambda_m^{a_m-b_m}] \sum_{l_1=0}^1 \sum_{l_2=0}^1 \dots \sum_{l_m=0}^1 (1 - l_1 - l_2 - \dots - l_m) \\ & \quad \times (\lambda_1)_{l_1} (\lambda_2)_{l_2} \dots (\lambda_m)_{l_m} e^{(A-a_1)\lambda_1} e^{(A-a_2)\lambda_2} \dots e^{(A-a_m)\lambda_m} \\ &= \left(\prod_{k=1}^m \frac{(A - a_k)^{a_k - b_k - 1}}{(a_k - b_k)!} \right) \sum_{l_1=0}^1 \sum_{l_2=0}^1 \dots \sum_{l_m=0}^1 (1 - l_1 - l_2 - \dots - l_m) \end{aligned} \quad (10)$$

$$\times \left(\prod_{j=1}^m (A - a_j)^{1-l_j} (a_j - b_j)_{l_j} \right). \quad (11)$$

Combining (10), (11), and (4), we obtain the main result:

Theorem 1

$$f(a_1, b_1; \dots; a_m, b_m) = \left(\prod_{k=1}^m \binom{a_k}{b_k} (A - a_k)^{a_k - b_k - 1} \right) \quad (12)$$

$$\times \sum_{l_1=0}^1 \sum_{l_2=0}^1 \dots \sum_{l_m=0}^1 (1 - l_1 - l_2 - \dots - l_m) \left(\prod_{j=1}^m (A - a_j)^{1-l_j} (a_j - b_j)_{l_j} \right). \quad (13)$$

For the case $m = 2$, formula (12), (13) specializes to the formula of Jin and Liu (1), and formula (12), (13) yields a closed formula for every fixed m . Note that for the case $m = 2$ we do not even have to evaluate the determinant in general, since for $m = 2$ simply

$$\det \begin{vmatrix} \delta_{ij} - (1 - \delta_{ij})\lambda_j \end{vmatrix} = 1 - \lambda_1 \lambda_2.$$

References

- [1] D. K. Faddeev, I. S. Sominskii, Problems in higher algebra, translated by J. L. Brenner, W. H. Freeman and Co., San Francisco-London, 1965.
- [2] I. P. Goulden and D. M. Jackson, Combinatorial Enumeration, John Wiley & Sons, New York, 1983.
- [3] Y. Jin and C. Liu, Enumeration for spanning forests of complete bipartite graphs, to appear in Ars Combinatoria.