

---

# MATH 532, 736I: MODERN GEOMETRY

## Test 2 Solutions

---

### Test 1 (1992):

#### Part I:

- (1) If  $A$ ,  $B$ , and  $C$  are not distinct, then the conclusion that  $A$ ,  $B$ , and  $C$  are collinear is clear. Suppose now that  $A$ ,  $B$ , and  $C$  are distinct. Recall that not all of  $x$ ,  $y$ , and  $z$  are zero. Suppose  $x \neq 0$  (a similar argument can be made if  $y \neq 0$  or  $z \neq 0$ ). From  $x\vec{A} + y\vec{B} + z\vec{C} = \vec{0}$ , we obtain  $\vec{A} = (-y/x)\vec{B} + (-z/x)\vec{C}$ . Let  $t = -z/x$ . Since  $x + y + z = 0$ ,

$$1 - t = 1 - \left(-\frac{z}{x}\right) = 1 + \frac{z}{x} = \frac{x+z}{x} = -\frac{y}{x}.$$

Hence,  $\vec{A} = (1-t)\vec{B} + t\vec{C}$ . By Theorem 1,  $\vec{A}$  is on  $\overleftrightarrow{BC}$ . Thus,  $A$ ,  $B$ , and  $C$  are collinear.

- (2) The square of the distance from  $N$  to  $M_A$  is  $(N - M_A)^2$  and the square of the distance from  $N$  to  $M_B$  is  $(N - M_B)^2$ . It therefore suffices to show that  $(N - M_A)^2 = (N - M_B)^2$ . Since  $\overleftrightarrow{CD}$  and  $\overleftrightarrow{BA}$  are perpendicular,  $(D - C)(A - B) = 0$ . We use that

$$\begin{aligned} D - C &= (A + B + C + D) - (B + C) - (A + C) \\ &= 4N - 2M_A - 2M_B = 2((N - M_A) + (N - M_B)). \end{aligned}$$

and  $A - B = (A + C) - (B + C) = 2M_B - 2M_A = 2((N - M_A) - (N - M_B))$

Since  $(D - C)(A - B) = 0$ , we deduce

$$0 = ((N - M_A) + (N - M_B))((N - M_A) - (N - M_B)) = (N - M_A)^2 - (N - M_B)^2.$$

Hence,  $(N - M_A)^2 = (N - M_B)^2$ .

#### Part II:

- (1) Not relevant to the current course.
- (2) Let  $P_A$  be the intersection of the altitude drawn from  $A$  to  $\overleftrightarrow{BC}$ . Let  $P_B$  be the intersection of the altitude drawn from  $B$  to  $\overleftrightarrow{AC}$ . Let  $P_C$  be the intersection of the altitude drawn from  $C$  to  $\overleftrightarrow{AB}$ . Let  $D$  be the intersection of  $\overleftrightarrow{AP_A}$  and  $\overleftrightarrow{BP_B}$ . Then  $\overleftrightarrow{AD}$  and  $\overleftrightarrow{BC}$  are perpendicular and  $\overleftrightarrow{BD}$  and  $\overleftrightarrow{AC}$  are perpendicular. Hence,  $(D - A)(C - B) = 0$  and  $(D - B)(C - A) = 0$ . Therefore,

$$0 = (D - A)(C - B) - (D - B)(C - A) = -AC - BD + AD + BC = (D - C)(A - B).$$

Thus,  $\overleftrightarrow{CD}$  and  $\overleftrightarrow{BA}$  are perpendicular. We deduce that the altitude from  $C$  also passes through  $D$  so that the three altitudes are concurrent.

- (3) In this problem,  $f = R_{\pi/2,A}T_AR_{\pi,A}$ . From the last page of this test (and as shown on homework from class)  $T_A = R_{\pi,A/2}R_{\pi,(0,0)}$ . Thus, we can view  $f$  as a product of four rotations with the sum of the angles in the rotations being  $7\pi/2$ . Since a rotation by  $7\pi/2$  is the same as a rotation by  $3\pi/2$ , we deduce that  $f = R_{3\pi/2,B}$ . Using  $A = (1, 3)$ , one writes  $R_{\pi/2,A}T_AR_{\pi,A}$  as a product of 3 matrices and obtains

$$R_{\pi/2,A}T_AR_{\pi,A} = \begin{pmatrix} 0 & 1 & -5 \\ -1 & 0 & 5 \\ 0 & 0 & 1 \end{pmatrix}.$$

(Note that the matrix above can be used to justify  $f = R_{3\pi/2,B}$  as well.) If  $B = (x, y)$ , then

$$R_{3\pi/2,B} = \begin{pmatrix} 0 & 1 & x - y \\ -1 & 0 & x + y \\ 0 & 0 & 1 \end{pmatrix}.$$

Hence,  $x - y = -5$  and  $x + y = 5$ . Solving, we deduce  $x = 0$  and  $y = 5$ . Thus,  $B = (0, 5)$ .

### Part III:

- (1) The vector  $B - A$  (going in the direction of  $\ell$ ) is parallel to the vector  $D - C$  (going in the direction of  $m$ ). Thus,  $B - A = k(D - C)$  for some constant  $k$ . Since  $E$  is the midpoint of  $\overline{AC}$  and  $F$  is the midpoint of  $\overline{BD}$ , we obtain

$$E = \frac{1}{2}(A + C) \quad \text{and} \quad F = \frac{1}{2}(B + D).$$

Thus,

$$F - E = \frac{1}{2}(B + D) - \frac{1}{2}(A + C) = \frac{1}{2}(B - A) + \frac{1}{2}(D - C) = \frac{k+1}{2}(D - C).$$

Thus,  $\overrightarrow{EF}$  is a constant times  $\overrightarrow{CD}$ . This constant is non-zero since  $E \neq F$ . It follows that  $\overrightarrow{EF}$  is parallel to  $\overrightarrow{CD}$ . Hence,  $\overrightarrow{EF}$  is parallel to both  $\ell$  and  $m$ .

- (2) You do not need to know this for Test 2.
- (3) **Proof.** By Theorem 1 (from the Information Page at the end of this test), there are real numbers  $k_1$ ,  $k_2$ , and  $k_3$  such that

$$X = (1 - k_1)A + k_1A' = (1 - k_2)B + k_2B' = (1 - k_3)C + k_3C'.$$

Next, we show that  $k_1 \neq k_2$ . Assume  $k_1 = k_2$ . Observe that  $k_1 \neq 0$  since otherwise we would have  $X = A = B$ , contradicting that  $A$  and  $B$  are distinct points. Also,  $k_1 \neq 1$  since otherwise we would have  $X = A' = B'$ , contradicting that  $A'$  and  $B'$  are distinct points. We get that

$$(1 - k_1)A - (1 - k_2)B = k_2B' - k_1A'$$

and that the vectors  $\overrightarrow{BA}$  and  $\overrightarrow{A'B'}$  either have the same direction or the exact opposite direction. This contradicts that the point  $X$  exists. Hence,  $k_1 \neq k_2$ . Thus,

$$\frac{1 - k_1}{k_2 - k_1}A + \frac{k_2 - 1}{k_2 - k_1}B = \frac{k_2}{k_2 - k_1}B' + \frac{-k_1}{k_2 - k_1}A'.$$

By Theorem 1 with  $t = (k_2 - 1)/(k_2 - k_1)$ , we see that the expression on the left above is a point on line  $\overleftrightarrow{AB}$ . By Theorem 1 with  $t = -k_1/(k_2 - k_1)$ , we see that the expression on the right above is a point on line  $\overleftrightarrow{A'B'}$ . Therefore, we get that

$$P = \frac{1 - k_1}{k_2 - k_1}A + \frac{k_2 - 1}{k_2 - k_1}B.$$

Hence,

$$(1) \quad (k_2 - k_1)P = (1 - k_1)A + (k_2 - 1)B.$$

Using that

$$(1 - k_2)B - (1 - k_3)C = k_3C' - k_2B',$$

we similarly obtain that  $k_2 \neq k_3$ , that

$$\frac{1 - k_2}{k_3 - k_2}B + \frac{k_3 - 1}{k_3 - k_2}C = \frac{k_3}{k_3 - k_2}C' + \frac{-k_2}{k_3 - k_2}B',$$

and that

$$(2) \quad (k_3 - k_2)Q = (1 - k_2)B + (k_3 - 1)C.$$

From

$$(1 - k_3)C - (1 - k_1)A = k_1A' - k_3C',$$

we similarly obtain that either

$$(3) \quad k_3 = k_1$$

or

$$(4) \quad \frac{1 - k_3}{k_1 - k_3}C + \frac{k_1 - 1}{k_1 - k_3}A = \frac{k_1}{k_1 - k_3}A' + \frac{-k_3}{k_1 - k_3}C'.$$

If (4) holds, then we could deduce that there is a point on both line  $\overleftrightarrow{AC}$  and line  $\overleftrightarrow{A'C'}$ , giving a contradiction. Thus, (3) must hold. We get from (1) and (2) that

$$(k_2 - k_1)P + (k_3 - k_2)Q = (1 - k_1)A + (k_3 - 1)C$$

so that

$$(k_2 - k_1)(P - Q) = (1 - k_1)(A - C).$$

Observe that  $P \neq Q$  since otherwise we would have that the points  $A$ ,  $B$ , and  $C$  are collinear, which isn't the case. Since  $k_1 \neq k_2$ , we obtain that the lines  $\overleftrightarrow{PQ}$  and  $\overleftrightarrow{AC}$  are parallel, completing the proof. ■

**Test 1 (1993):****Part I:**

- (1) If  $A = B$ , then take  $x = 1$ ,  $y = -1$ , and  $z = 0$ . Suppose now that  $A \neq B$ . By Theorem 1, there is a real number  $t$  such that  $C = (1 - t)A + tB$ . Let  $x = 1 - t$ ,  $y = t$ , and  $z = -1$ . Then  $x$ ,  $y$ , and  $z$  are not all 0,  $x + y + z = 0$ , and  $xA + yB + zC = \vec{0}$ .
- (2) See Problem (2) of the 1992 test.
- (3) One answer is:  $\triangle AA'Z$  and  $\triangle BB'X$  are perspective from point  $Y$ .

**Part II:**

- (1) It suffices to show that  $G = \frac{1}{2}(C + F)$ . The given information implies

$$D = \frac{B + C}{2}, \quad E = \frac{A + C}{2}, \quad F = \frac{A + B}{2}, \quad \text{and} \quad G = \frac{D + E}{2}.$$

Hence,

$$G = \frac{1}{2}(D + E) = \frac{1}{2}\left(\frac{B + C}{2} + \frac{A + C}{2}\right) = \frac{1}{2}\left(C + \frac{A + B}{2}\right) = \frac{1}{2}(C + F),$$

as desired.

- (2) From the given information,  $f = R_{\pi/2, (1,1)}R_{\pi, (1,0)}R_{\pi/2, (0,0)}$ . Since the sum of the angles ( $\pi/2$ ,  $\pi$ , and  $\pi/2$ ) is  $2\pi$  (an integer times  $2\pi$ ), we deduce that  $f$  is a translation. In other words,  $f = T_B$ . To determine  $B$ , one can compute  $f$  by multiplying matrices. However, it is probably easier to take a point and see what  $f$  does to it (where it is mapped under  $f$ ). Consider  $(x, y) = (0, 0)$ . Rotating this point about  $(0, 0)$  by  $\pi/2$  does not change it. Now, rotating the point about  $(1, 0)$  by  $\pi$  moves it to  $(2, 0)$ . Finally, rotating this point about  $(1, 1)$  by  $\pi/2$  moves it to  $(2, 2)$ . Thus,  $f$  takes  $(0, 0)$  to  $(2, 2)$ . It follows that  $B = (2, 2)$ .
- (3) You do not need to know this for Test 2.
- (4) Recall in class that we discussed how one could obtain the point  $C$  where  $R_{\alpha+\beta, C} = R_{\beta, B}R_{\alpha, A}$ . The points  $A$ ,  $B$ , and  $C$  form a triangle with  $\angle BAC = \alpha/2$  and  $\angle ABC = \beta/2$ . Since  $\alpha + \beta = \pi$  in this problem, the sum of the measures of  $\angle BAC$  and  $\angle ABC$  is  $\pi/2$ . Thus,  $\triangle ABC$  is a right triangle with  $\angle ACB = \pi/2$ . Recall that we showed in class that in this situation,  $C$  is on the circle having diameter  $\overline{AB}$ . Since the center of this circle is the midpoint of  $\overline{AB}$ ,  $C$  is on the circle centered at  $\frac{1}{2}(A + B)$  passing through  $A$  and  $B$ .

**Test 1 (1994):**

- (1) See Problem 1 on the 1992 test. Note that the theorems are numbered differently.
- (2) Observe that

$$2N - M_A - Q_A = \frac{A + B + C + D}{2} - \frac{B + C}{2} - \frac{A + D}{2} = \vec{0}.$$

Thus,

$$\begin{aligned} 0 &= (2N - M_A - Q_A)(M_A - Q_A) = 2NM_A - M_A^2 - 2NQ_A + Q_A^2 \\ &= -N^2 + 2NM_A - M_A^2 + N^2 - 2NQ_A + Q_A^2 = -(N - M_A)^2 + (N - Q_A)^2. \end{aligned}$$

Thus,  $(N - M_A)^2 = (N - Q_A)^2$ , and we deduce that the distance from  $N$  to  $M_A$  is the same as the distance from  $N$  to  $Q_A$ .

- (3) (a)  $(3, -2)$
- (b) 4
- (c)  $(4, 4)$  (you should be able to do this with and without matrices)
- (d)  $(1, 4)$  (you should be able to do this with and without matrices)
- (4) One answer is:  $\triangle AZA'$  and  $\triangle BXB'$  are perspective from point  $Y$ . There are other answers. Another one that was suggested is:  $\triangle A'YA$  and  $\triangle C'XC$  are perspective from point  $Z$ .
- (5) Observe that  $a^2 = (B - A)^2$ ,  $b^2 = (C - B)^2$ , and  $c^2 = (C - A)^2$ . Since  $a^2 + b^2 = c^2$ , we obtain

$$(B - A)^2 + (C - B)^2 = (C - A)^2$$

so that

$$B^2 - 2AB + A^2 + C^2 - 2BC + B^2 = C^2 - 2AC + A^2.$$

Rearranging, we obtain  $2B^2 - 2AB - 2BC + 2AC = 0$ . Dividing by 2, we obtain  $0 = B^2 - AB - BC + AC = (A - B)(C - B)$ . Therefore,  $\vec{BA}$  and  $\vec{BC}$  are perpendicular and, hence,  $\angle ABC$  is a right angle.

- (6) You do not need to know this for Test 2.
- (7) Since  $f$  is three successive rotations with the angles of these rotations summing to  $8\pi/3$ ,  $f = R_{8\pi/3, (x, y)} = R_{\alpha, D}$  where  $\alpha = 2\pi/3$  and  $D = (x, y)$  for some numbers  $x$  and  $y$ . We need to determine  $x$  and  $y$ . Since  $f(1, 1) = (5, 7)$ , we deduce that

$$\begin{pmatrix} 5 \\ 7 \\ 1 \end{pmatrix} = R_{2\pi/3, (x, y)} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} -1/2 & -\sqrt{3}/2 & (3x + \sqrt{3}y)/2 \\ \sqrt{3}/2 & -1/2 & (-\sqrt{3}x + 3y)/2 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$$

Thus,  $10 = -1 - \sqrt{3} + 3x + \sqrt{3}y$  and  $14 = \sqrt{3} - 1 - \sqrt{3}x + 3y$ .

Rewriting these, we obtain  $3x + \sqrt{3}y = 11 + \sqrt{3}$  and  $\sqrt{3}x - 3y = -15 + \sqrt{3}$ .

Multiplying the first of these by  $\sqrt{3}$  and adding the result to the second, we deduce  $4\sqrt{3}x = 12\sqrt{3} - 12$  so that  $x = 3 - \sqrt{3}$ . Substituting, we obtain (after a little work)  $y = 4 + (2\sqrt{3}/3)$ .

(8) **Proof:** Since  $\overleftrightarrow{AA'}$ ,  $\overleftrightarrow{BB'}$ , and  $\overleftrightarrow{CC'}$  are parallel, there are real numbers  $k_1$  and  $k_2$  such that

$$A' - A = k_1(B' - B) = k_2(C' - C).$$

We get that

$$A - k_1B = A' - k_1B'.$$

We first explain why  $k_1 \neq 1$ . Assume  $k_1 = 1$ . Then  $A - B = A' - B'$  so that  $\overleftrightarrow{AB}$  and  $\overleftrightarrow{A'B'}$  are parallel. This contradicts that  $P$  exists. Therefore,  $k_1 \neq 1$ . Hence,

$$\left(\frac{1}{1-k_1}\right)A + \left(\frac{-k_1}{1-k_1}\right)B = \left(\frac{1}{1-k_1}\right)A' + \left(\frac{-k_1}{1-k_1}\right)B'.$$

By Theorem 1 (from the last page of this exam) with  $t = -k_1/(1-k_1)$ , we see that the expression on the left above is a point on line  $\overleftrightarrow{AB}$  and that the expression on the right above is a point on line  $\overleftrightarrow{A'B'}$ . Therefore, we get that

$$(1) \quad (1-k_1)P = A - k_1B.$$

Similarly, from  $k_1B - k_2C = k_1B' - k_2C'$ , we deduce that

$$(2) \quad (k_1 - k_2)Q = k_1B - k_2C.$$

Also, from  $k_2C - A = k_2C' - A'$ , we deduce that

$$(3) \quad (k_2 - 1)R = k_2C - A.$$

Therefore, from (1), (2), and (3),

$$(1-k_1)P + (k_1 - k_2)Q + (k_2 - 1)R = \vec{0}.$$

The result follows from Theorem 3 (on the last page of this test).

**Test 1 (1995):**

- (1) See Problem 1 on the 1993 test. Note that the theorems are numbered differently.
- (2) The square of the distance from  $N$  to  $M_A$  is  $(N - M_A)^2$  and the square of the distance from  $N$  to  $M_C$  is  $(N - M_C)^2$ . It therefore suffices to show that  $(N - M_A)^2 = (N - M_C)^2$ . Since  $\overleftrightarrow{BD}$  and  $\overleftrightarrow{CA}$  are perpendicular,  $(D - B)(A - C) = 0$ . We use that

$$\begin{aligned} D - B &= (A + B + C + D) - (B + C) - (A + B) \\ &= 4N - 2M_A - 2M_C = 2((N - M_A) + (N - M_C)). \end{aligned}$$

and  $A - C = (A + B) - (B + C) = 2M_C - 2M_A = 2((N - M_A) - (N - M_C))$

Since  $(D - B)(A - C) = 0$ , we deduce

$$0 = ((N - M_A) + (N - M_C))((N - M_A) - (N - M_C)) = (N - M_A)^2 - (N - M_C)^2.$$

Hence,  $(N - M_A)^2 = (N - M_C)^2$ .

- (3) (a)  $(-2, 0)$   
 (b) 1  
 (c)  $(12, 6)$
- (4) Observe that  $a^2 = (B - A)^2$ ,  $b^2 = (C - B)^2$ , and  $c^2 = (C - A)^2$ . Since  $a^2 + b^2 = c^2$ , we obtain

$$(B - A)^2 + (C - B)^2 = (C - A)^2$$

so that

$$B^2 - 2AB + A^2 + C^2 - 2BC + B^2 = C^2 - 2AC + A^2.$$

Rearranging, we obtain  $2B^2 - 2AB - 2BC + 2AC = 0$ . Dividing by 2, we obtain  $0 = B^2 - AB - BC + AC = (A - B)(C - B)$ . Therefore,  $\overleftrightarrow{BA}$  and  $\overleftrightarrow{BC}$  are perpendicular and, hence,  $\angle ABC$  is a right angle.

- (5) One answer is:  $\triangle TB'B$  and  $\triangle SA'A$  are perspective from point  $R$ .
- (6) Let  $D$  be the midpoint of  $\overleftrightarrow{BC}$ . Then  $D = (B + C)/2$ . Since the distance from  $A$  to  $B$  is equal to the distance from  $A$  to  $C$ , we obtain  $(A - B)^2 = (A - C)^2$ . Thus,

$$\begin{aligned} 0 &= (A - B)^2 - (A - C)^2 = (2A - B - C)(C - A) \\ &= 2\left(A - \frac{B + C}{2}\right)(C - A) = 2(A - D)(C - A) \end{aligned}$$

(where the second equality follows by considering the factorization of the difference of two squares). It follows that  $\overleftrightarrow{DA}$  and  $\overleftrightarrow{AC}$  are perpendicular. In other words, the line passing through  $A$  and the midpoint of  $\overleftrightarrow{BC}$  is perpendicular to line  $\overleftrightarrow{BC}$ .

(7) **Proof:** Since  $\overleftrightarrow{AA'}$ ,  $\overleftrightarrow{BB'}$ , and  $\overleftrightarrow{CC'}$  are parallel, there are real numbers  $k_1$  and  $k_2$  such that

$$A' - A = k_1(B' - B) = k_2(C' - C).$$

We get that

$$A - k_1B = A' - k_1B'.$$

We first explain why  $k_1 \neq 1$ . Assume  $k_1 = 1$ . Then  $A - B = A' - B'$  so that  $\overleftrightarrow{AB}$  and  $\overleftrightarrow{A'B'}$  are parallel. This contradicts that  $P$  exists. Therefore,  $k_1 \neq 1$ . Hence,

$$\left(\frac{1}{1-k_1}\right)A + \left(\frac{-k_1}{1-k_1}\right)B = \left(\frac{1}{1-k_1}\right)A' + \left(\frac{-k_1}{1-k_1}\right)B'.$$

By Theorem 1 (from the last page of this exam) with  $t = -k_1/(1-k_1)$ , we see that the expression on the left above is a point on line  $\overleftrightarrow{AB}$  and that the expression on the right above is a point on line  $\overleftrightarrow{A'B'}$ . Therefore, we get that

$$(1) \quad (1-k_1)P = A - k_1B.$$

Similarly, from  $k_1B - k_2C = k_1B' - k_2C'$ , we deduce that

$$(2) \quad (k_1 - k_2)Q = k_1B - k_2C.$$

Using that

$$A - k_2C = A' - k_2C'$$

and that  $\overleftrightarrow{AC}$  and  $\overleftrightarrow{A'C'}$  are parallel (so  $R$  is a point at “infinity”), we obtain

$$(3) \quad k_2 = 1.$$

From (1) and (2), we obtain

$$(4) \quad (1-k_1)P + (k_1 - k_2)Q = A - k_2C.$$

Using (3), we can rewrite (4) in the form

$$(1-k_1) \times (P - Q) = A - C.$$

Recall that  $k_1 \neq 1$ . Therefore, the line  $\overleftrightarrow{AC}$  is parallel to the line  $\overleftrightarrow{PQ}$ .