

MARK BOX		
PROBLEM	POINTS	
1 a - y	25	
2 a - o	15	
3	3	
4 a - c	5	
5	5	
6	5	
7	5	
8	5	
9	5	
10	5	
11	5	
12	5	
13	5	
14	5	
15	5	
16	5	
POSSIBLE	108	

NAME: \_\_\_\_\_

please check the box of your section

Section 001 (MW 9:05 am)

or

Section 002 (MW 10:10 am)

**INSTRUCTIONS:**

- (1) To receive credit you must:
  - (a) **work in a logical fashion, show all your work, indicate your reasoning; no credit will be given for an answer that *just appears*; such explanations help with partial credit**
  - (b) if a line/box is provided, then:
    - show you work BELOW the line/box
    - put your answer on/in the line/box
  - (c) if no such line/box is provided, then box your answer
- (2) The MARK BOX indicates the problems along with their points. Check that your copy of the exam has all of the problems.
- (3) You may **not** use a calculator, books, personal notes.
- (4) During this exam, do not leave your seat. If you have a question, raise your hand. When you finish: turn your exam over, put your pencil down, and raise your hand.
- (5) This exam covers (from *Calculus* by Anton, Bivens, Davis 8<sup>th</sup> ed.):  
 $S$  7.1 - 7.4, 7.6, 7.7, 8.1 - 8.5, 8.8, 10.1-10.10, 11.1 - 11.3. .

**1. Fill in the blanks (each worth 1 point).**

**1a.**  $\int \frac{du}{u} = \underline{\hspace{2cm}} |u| + C$

**1b.** If  $a$  is a constant and  $a > 0$  but  $a \neq 1$ , then  $\int a^u du = \underline{\hspace{2cm}} + C$

**1c.**  $\int \cos u du = \underline{\hspace{2cm}} + C$

**1d.**  $\int \sec^2 u du = \underline{\hspace{2cm}} + C$

**1e.**  $\int \sec u \tan u du = \underline{\hspace{2cm}} + C$

**1f.**  $\int \sin u du = \underline{\hspace{2cm}} + C$

**1g.**  $\int \csc^2 u du = \underline{\hspace{2cm}} + C$

**1h.**  $\int \csc u \cot u du = \underline{\hspace{2cm}} + C$

**1i.**  $\int \tan u du = \underline{\hspace{2cm}} + C$

**1j.**  $\int \cot u du = \underline{\hspace{2cm}} + C$

**1k.**  $\int \sec u du = \underline{\hspace{2cm}} + C$

**1l.**  $\int \csc u du = \underline{\hspace{2cm}} + C$

**1m.** If  $a$  is a constant and  $a > 0$  then  $\int \frac{1}{\sqrt{a^2-u^2}} du = \underline{\hspace{2cm}} + C$

**1n.** If  $a$  is a constant and  $a > 0$  then  $\int \frac{1}{a^2+u^2} du = \underline{\hspace{2cm}} + C$

**1o.** If  $a$  is a constant and  $a > 0$  then  $\int \frac{1}{u\sqrt{u^2-a^2}} du = \underline{\hspace{2cm}} + C$

**1p.** Partial Fraction Decomposition. If one wants to integrate  $\frac{f(x)}{g(x)}$  where  $f$  and  $g$  are polynomials and  $[\text{degree of } f] \geq [\text{degree of } g]$ , then one must first do  $\underline{\hspace{2cm}}$

**1q.** Integration by parts formula:  $\int u dv = \underline{\hspace{2cm}}$

**1r.** Trig substitution: (recall that the *integrand* is the function you are integrating)  
if the integrand involves  $a^2-u^2$ , then one makes the substitution  $u = \underline{\hspace{2cm}}$

**1s.** Trig substitution:  
if the integrand involves  $a^2+u^2$ , then one makes the substitution  $u = \underline{\hspace{2cm}}$

**1t.** Trig substitution:  
if the integrand involves  $u^2-a^2$ , then one makes the substitution  $u = \underline{\hspace{2cm}}$

**1u.** trig formula ... your answer should involve trig functions of  $\theta$ , and not of  $2\theta$ :  $\sin(2\theta) = \underline{\hspace{2cm}}$ .

**1v.** trig formula ...  $\cos(2\theta)$  should appear in the numerator:  $\cos^2(\theta) = \frac{\underline{\hspace{2cm}}}{2}$ .

**1w.** trig formula ...  $\cos(2\theta)$  should appear in the numerator:  $\sin^2(\theta) = \frac{\underline{\hspace{2cm}}}{2}$ .

**1x.** trig formula ... since  $\cos^2 \theta + \sin^2 \theta = 1$ , we know that the corresponding relationship between tangent (i.e.,  $\tan$ ) and secant (i.e.,  $\sec$ ) is  $\underline{\hspace{2cm}}$ .

**1y.**  $\arctan\left(\frac{-1}{\sqrt{3}}\right) = \underline{\hspace{2cm}}$  RADIANS. (your answer should be an angle)

2. Fill-in-the blanks/boxes. All series  $\sum$  are understood to be  $\sum_{n=1}^{\infty}$ .

**Hint: I do NOT want to see the words absolute nor conditional on this page!**

**2a. Sequences** Let  $-\infty < r < \infty$ . (Fill-in-the blanks with *exists* or *does not exist*, i.e. *DNE*)

- If  $|r| < 1$ , then  $\lim_{n \rightarrow \infty} r^n$  \_\_\_\_\_
- If  $|r| > 1$ , then  $\lim_{n \rightarrow \infty} r^n$  \_\_\_\_\_
- If  $r = 1$ , then  $\lim_{n \rightarrow \infty} r^n$  \_\_\_\_\_
- If  $r = -1$ , then  $\lim_{n \rightarrow \infty} r^n$  \_\_\_\_\_

**2b. Geometric Series** where  $-\infty < r < \infty$ . The series  $\sum r^n$

- converges if and only if  $|r|$  \_\_\_\_\_
- diverges if and only if  $|r|$  \_\_\_\_\_

**2c.  $p$ -series** where  $0 < p < \infty$ . The series  $\sum \frac{1}{n^p}$

- converges if and only if  $p$  \_\_\_\_\_
- diverges if and only if  $p$  \_\_\_\_\_

**2d. Integral Test** for a positive-termed series  $\sum a_n$  where  $a_n \geq 0$ .

Let  $f: [1, \infty) \rightarrow \mathbb{R}$  be so that

- $a_n = f(\text{_____})$  for each  $n \in \mathbb{N}$
- $f$  is a \_\_\_\_\_ function
- $f$  is a \_\_\_\_\_ function
- $f$  is a \_\_\_\_\_ function .

Then  $\sum a_n$  converges if and only if \_\_\_\_\_ converges.

**2e. Comparison Test** for a positive-termed series  $\sum a_n$  where  $a_n \geq 0$ .

- If  $0 \leq a_n \leq b_n$  for all  $n \in \mathbb{N}$  and  $\sum b_n$  \_\_\_\_\_, then  $\sum a_n$  \_\_\_\_\_.
- If  $0 \leq b_n \leq a_n$  for all  $n \in \mathbb{N}$  and  $\sum b_n$  \_\_\_\_\_, then  $\sum a_n$  \_\_\_\_\_.

**2f. Limit Comparison Test** for a positive-termed series  $\sum a_n$  where  $a_n \geq 0$ .

Let  $b_n > 0$  and  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = L$ .

If \_\_\_\_\_  $< L <$  \_\_\_\_\_, then  $\sum a_n$  converges if and only if \_\_\_\_\_ .

**2g. Ratio and Root Tests** for a positive-termed series  $\sum a_n$  where  $a_n \geq 0$ .

Let  $\rho = \lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n}$  or  $\rho = \lim_{n \rightarrow \infty} (a_n)^{\frac{1}{n}}$ .

- If  $\rho$  \_\_\_\_\_ then  $\sum a_n$  converges.
- If  $\rho$  \_\_\_\_\_ then  $\sum a_n$  diverges.
- If  $\rho$  \_\_\_\_\_ then the test is inconclusive.

**2h. Alternating Series Test** for an alternating series  $\sum (-1)^n a_n$  where  $a_n > 0$  for each  $n \in \mathbb{N}$ .

If

- $a_n$  \_\_\_\_\_  $a_{n+1}$  for each  $n \in \mathbb{N}$
- $\lim_{n \rightarrow \infty} a_n =$  \_\_\_\_\_

then  $\sum (-1)^n a_n$  \_\_\_\_\_

**2i.  $n^{\text{th}}$ -term test** for an arbitrary series  $\sum a_n$ .

If  $\lim_{n \rightarrow \infty} a_n \neq 0$  or  $\lim_{n \rightarrow \infty} a_n$  does not exist, then  $\sum a_n$  \_\_\_\_\_ .

2j. By definition, for an arbitrary series  $\sum a_n$ , (fill in the blanks with converges or diverges).

- $\sum a_n$  is absolutely convergent if and only if  $\sum |a_n|$  \_\_\_\_\_
- $\sum a_n$  is conditionally convergent if and only if  $\sum a_n$  \_\_\_\_\_ and  $\sum |a_n|$  \_\_\_\_\_
- $\sum a_n$  is divergent if and only if  $\sum a_n$  \_\_\_\_\_

2k. Consider a **function**  $y = f(x)$  where  $f: [1, \infty) \rightarrow \mathbb{R}$ .

Next consider the corresponding **sequence**  $\{a_n\}_{n=1}^{\infty}$  where  $a_n \stackrel{\text{def.}}{=} f(n)$ .

- If the limit of the **function**  $y = f(x)$  as  $x \rightarrow \infty$  is  $L$ ,

then the limit of the corresponding **sequence**  $\{a_n\}_{n=1}^{\infty}$  as  $n \rightarrow \infty$  is \_\_\_\_\_.

- If  $\lim_{n \rightarrow \infty} a_n = L$ , is it necessarily true that  $\lim_{x \rightarrow \infty} f(x) = L$ ? Circle: **Yes** or **No**

for 2l – 2o

Let  $y = f(x)$  be a function with derivatives of all orders in an interval  $I$  containing  $x_0$ .

Let  $y = p_N(x)$  be the  $N^{\text{th}}$ -order Taylor polynomial of  $y = f(x)$  about  $x_0$ .

Let  $y = R_N(x)$  be the  $N^{\text{th}}$ -order Taylor remainder of  $y = f(x)$  about  $x_0$ .

Let  $y = p_{\infty}(x)$  be the Taylor series of  $y = f(x)$  about  $x_0$ .

2l. In open form (i.e., with ... and without a  $\sum$ -sign)

$$p_N(x) =$$

2m. In closed form (i.e., with a  $\sum$ -sign and without ...)

$$p_N(x) =$$


2n. In closed form (i.e., with a  $\sum$ -sign and without ...)

$$p_{\infty}(x) =$$


2o. We know that  $f(x) = p_N(x) + R_N(x)$ . Taylor's BIG Theorem tells us that, for each  $x \in I$ ,

$$R_N(x) =$$

for some  $c$  between




and



.

3. Fill in the 3 blank lines (with absolutely convergent, conditional convergent, or divergent) on the following FLOW CHART for class used to determine if a series  $\sum_{n=17}^{\infty} a_n$  is: absolutely convergent, conditional convergent, or divergent.



4. Let  $R$  be the region in the  $x - y$ -plane that is enclosed by

$$y = x \quad \text{and} \quad y = 4x + 1 \quad \text{and} \quad x = 0 \quad \text{and} \quad x = 1 .$$

In each of problems **4b** and **4c**:

- $R$  will be revolved around some line to create a solid of revolution
- using either the disk, washer, or shell method, express the volume  $V$  of the resulting solid of revolution as **one integral** (and NOT as 2 or more integrals).
- In the space provided **below** each problem, make some *good enough sketch* (does not have to be too fancy) to indicate (i.e., help justify) your thinking/reasoning behind your solution
- you do not have to do lots of algebra to your integrand
- you do not have to integrate your integral.

**4a.** Make a sketch of  $R$  below and label important points. Also, in your sketch of  $R$  below, draw in a typical rectangle (should it be horizontal or vertical?) that would be used to express the area of  $R$  as precisely 1 integral (and not 2 integrals).

**4b.** The volume  $V$  of the solid obtained by revolving the region  $R$  about the  $x$ -axis is

$V =$
-------

Hint  $(r_{\text{big}}^2 - r_{\text{little}}^2) \neq (r_{\text{big}} - r_{\text{little}})^2$

continued next page  
 $\Rightarrow$

4c. The volume  $V$  of the solid obtained by revolving the region  $R$  about the line  $x = 2$  is

$V =$

Hint: the line  $x = 2$  is a vertical line (it is NOT a horizontal line).

5.

$$\int \sec^3 x \tan^3 x \, dx = \qquad \qquad \qquad +C$$

Remark: box your substitution box for more partial credit.

Recall: you can check your answer via differentiation (if you have time).

6.

$$\int \frac{dx}{(4+x^2)^2} =$$

$+C$

Remark: box your substitution box for more partial credit.

Recall: you can check your answer via differentiation (if you have time).

Hint: useful might be problems: 1r – 1w.

7.

$$\int \frac{x^3 + x^2 + 2x + 1}{x^4 + 2x^2 + 1} dx =$$

$+C$

Remark: box your substitution box for more partial credit.

Recall: you can check your answer via differentiation (if you have time).

Hint:  $x^4 + 2x^2 + 1 = (x^2 + 1)^2$ , which is a irreducible quadratic squared.

8.

$$\int x^3 e^{x^2} dx =$$

$+C$

Remark: box your substitution box for more partial credit.

Recall: you can check your answer via differentiation (if you have time).

Hint: one lesson from parts was to look for a  $dv$  that is easy to integrate and thus get the  $v$ .

9.

$$\int \sin(\ln x) dx =$$

$+C$

Remark: box your substitution box for more partial credit.

Recall: you can check your answer via differentiation (if you have time).

Hint: one lesson from parts was that if the integrand is easy to differentiate but hard to integrate, then let  $u$  be in integrand. Another lesson from parts was the bring to the other side idea.

10. Check the correct box and then indicate your reasoning below. Specifically specify what test(s) you are using. A correctly checked box without appropriate explanation will receive no points.

$$\sum_{n=17}^{\infty} (-1)^n \frac{1}{\sqrt{n^2 - 1}}$$

- absolutely convergent
- conditionally convergent
- divergent

11. Check the correct box and then indicate your reasoning below. Specifically specify what test(s) you are using. A correctly checked box without appropriate explanation will receive no points.

$$\sum_{n=8}^{\infty} (-1)^n \frac{n!}{(2n-1)!}$$

absolutely convergent  
 conditionally convergent  
 divergent

But before you get started .... let

$$a_n = \frac{n!}{(2n-1)!} .$$

Then  $a_{n+1} =$

Next, simplify  $\frac{a_{n+1}}{a_n}$  so that it has NO factorial sign (that is a ! sign) in it.

$$\frac{a_{n+1}}{a_n} =$$

Ok, now you should be ready to finish off the problem and check the correct box above.

12. Consider the formal power series

$$\sum_{n=1}^{\infty} \frac{5^n (x-3)^n}{n^2}$$

Hint:  $\left| \frac{5^{n+1}(x-3)^{n+1}}{(n+1)^2} \frac{n^2}{5^n(x-3)^n} \right| = \frac{5^{n+1} |x-3|^{n+1}}{5^n |x-3|^n} \frac{n^2}{(n+1)^2} = 5 |x-3| \left( \frac{n}{n+1} \right)^2$ .

The center is  $x_0 =$  \_\_\_\_\_ and the radius of convergence is  $R =$  \_\_\_\_\_.

As we did in class, make a number line indicating where the power series is: absolutely convergent, conditionally convergent, and divergent. Indicate your reasoning and specifically specify what test(s) you are using. Don't forget to check the endpoints, if there are any.



13. Consider the formal power series

$$\sum_{n=1}^{\infty} \frac{n!}{n^3} (x-2)^n$$

Hint: do the same kind of calculation as done in the hint for the previous problem.

The center is  $x_0 =$  \_\_\_\_\_ and the radius of convergence is  $R =$  \_\_\_\_\_.

As we did in class, make a number line indicating where the power series is: absolutely convergent, conditionally convergent, and divergent. Indicate your reasoning and specifically specify what test(s) you are using. Don't forget to check the endpoints, if there are any.



14. Consider the point, in **polar coordinates**,

$$P = (r, \theta) = \left(2, \frac{\pi}{6}\right) .$$

In **cartesian coordinates**, the point  $P$  is given by

$$P = (x, y) = ( \underline{\hspace{2cm}} , \underline{\hspace{2cm}} ) .$$

Below graph, and CLEARLY label, the following points.

$$P = \left(2, \frac{\pi}{6}\right)$$

$$Q = \left(-2, \frac{\pi}{6}\right)$$

$$R = \left(2, -\frac{\pi}{6}\right)$$

$$S = \left(-2, -\frac{\pi}{6}\right) .$$

15. Consider the curve in polar coordinate

$$r = 3 \sin(2\theta) .$$

15a. The period of  $r = 3 \sin(2\theta)$  is \_\_\_\_\_.

15b.  $\frac{\text{the period of } r = 3 \sin(2\theta)}{4} =$  \_\_\_\_\_

15c. Make a chart, as we did in class, to help you graph  $r = 3 \sin(2\theta)$ .

15d. Graph  $r = 3 \sin(2\theta)$ .

16. Express the area enclosed by  $r = 3 \sin(2\theta)$  as an integral with respect to  $\theta$   
(ok ... with respect to  $\theta$  means a  $d\theta$  in there).  
(You do not have to evaluate this integral.)

area =