

MARK BOX		
PROBLEM	POINTS	
1 a–y	25	
2	10	
3	10	
4	10	
5	10	
6	10	
7	10	
take home	10	
Extra Credit	5	
%	100	

**NAME** (legibly printed): \_\_\_\_\_

**class PIN:** \_\_\_\_\_

(\*) Extra Credit: 5 point for knowing your PIN number.

**INSTRUCTIONS:**

- (1) To receive credit you must:
  - (a) **work in a logical fashion, show all your work, indicate your reasoning;**  
no credit will be given for an answer that *just appears*;  
 such explanations help with partial credit
  - (b) if a line/box is provided, then:
    - show your work BELOW the line/box
    - put your answer on/in the line/box
  - (c) if no such line/box is provided, then box your answer
- (2) The MARK BOX indicates the problems along with their points.  
 Check that your copy of the exam has all of the problems.
- (3) You may **not** use a calculator, books, personal notes.
- (4) During this exam, do not leave your seat. If you have a question, raise your hand. When you finish: turn your exam over, put your pencil down, and raise your hand.
- (5) This exam covers (from *Calculus* by Anton, Bivens, Davis 8<sup>th</sup> ed.):  
 Sections 8.1, 8.2, 8.3, 8.4, 8.5, 8.7, 8.8. .

**Problem Inspiration:** If I told you here, you would know what method to use. So see the solution key, which will be available from the course homepage shortly after the exam.

**Hints:**

- (1) **You can check your answers to the indefinite integrals by differentiating.**
- (2) **+ C**
- (3) **For more partial credit, box your  $u - du$  substitutions.**

**1. Fill in the blanks (each worth 1 point).**

**1a.**  $\int \frac{du}{u} = \underline{\hspace{2cm}} |u| + C$

**1b.** If  $a$  is a constant and  $a > 0$  but  $a \neq 1$ , then  $\int a^u du = \underline{\hspace{2cm}} + C$

**1c.**  $\int \cos u du = \underline{\hspace{2cm}} + C$

**1d.**  $\int \sec^2 u du = \underline{\hspace{2cm}} + C$

**1e.**  $\int \sec u \tan u du = \underline{\hspace{2cm}} + C$

**1f.**  $\int \sin u du = \underline{\hspace{2cm}} + C$

**1g.**  $\int \csc^2 u du = \underline{\hspace{2cm}} + C$

**1h.**  $\int \csc u \cot u du = \underline{\hspace{2cm}} + C$

**1i.**  $\int \tan u du = \underline{\hspace{2cm}} + C$

**1j.**  $\int \cot u du = \underline{\hspace{2cm}} + C$

**1k.**  $\int \sec u du = \underline{\hspace{2cm}} + C$

**1l.**  $\int \csc u du = \underline{\hspace{2cm}} + C$

**1m.** If  $a$  is a constant and  $a > 0$  then  $\int \frac{1}{\sqrt{a^2-u^2}} du = \underline{\hspace{2cm}} + C$

**1n.** If  $a$  is a constant and  $a > 0$  then  $\int \frac{1}{a^2+u^2} du = \underline{\hspace{2cm}} + C$

**1o.** If  $a$  is a constant and  $a > 0$  then  $\int \frac{1}{u\sqrt{u^2-a^2}} du = \underline{\hspace{2cm}} + C$

**1p.** Partial Fraction Decomposition. If one wants to integrate  $\frac{f(x)}{g(x)}$  where  $f$  and  $g$  are polynomials

and  $[\text{degree of } f] \geq [\text{degree of } g]$ , then one must first do  $\underline{\hspace{2cm}}$

**1q.** Integration by parts formula:  $\int u dv = \underline{\hspace{2cm}}$

**1r.** Trig substitution: (recall that the *integrand* is the function you are integrating)  
if the integrand involves  $a^2-u^2$ , then one makes the substitution  $u = \underline{\hspace{2cm}}$

**1s.** Trig substitution:  
if the integrand involves  $a^2+u^2$ , then one makes the substitution  $u = \underline{\hspace{2cm}}$

**1t.** Trig substitution:  
if the integrand involves  $u^2-a^2$ , then one makes the substitution  $u = \underline{\hspace{2cm}}$

**1u.** trig formula ... your answer should involve trig functions of  $\theta$ , and not of  $2\theta$ :  $\sin(2\theta) = \underline{\hspace{2cm}}$ .

**1v.** trig formula ...  $\cos(2\theta)$  should appear in the numerator:  $\cos^2(\theta) = \frac{\underline{\hspace{2cm}}}{2}$ .

**1w.** trig formula ...  $\cos(2\theta)$  should appear in the numerator:  $\sin^2(\theta) = \frac{\underline{\hspace{2cm}}}{2}$ .

**1x.** trig formula ... since  $\cos^2 \theta + \sin^2 \theta = 1$ , we know that the corresponding relationship between tangent (i.e., tan) and secant (i.e., sec) is  $\underline{\hspace{2cm}}$ .

**1y.**  $\arctan(-\sqrt{3}) = \underline{\hspace{2cm}}$  **RADIANS**. (your answer should be an angle)

2.

$$\int (\sec^3 x) (\tan^3 x) dx =$$

+ C

3.

$$\int \frac{1}{(4+x^2)^2} dx =$$

+ C

4.

$$\int \frac{x^3 + x^2 + 2x + 1}{x^4 + 2x^2 + 1} dx = \quad + C$$

Hint:  $x^4 + 2x^2 + 1 = (x^2 + 1)^2$  .

5.

$$\int x^3 e^{x^2} dx =$$

+ C

6.

$$\int \sin(\ln x) dx =$$

+ C

Hint: bring to the other side idea.

7.

$$\int_0^e \frac{dx}{x-2} =$$

HINT:  $e \approx 2.7 > 2$