

MARK BOX		
PROBLEM	POINTS	
1	25	
2	5	
3	10	
4	10	
5	10	
6	10	
7	10	
8	10	
take home	10	
%	100	

NAME: \_\_\_\_\_

class PIN: \_\_\_\_\_

**INSTRUCTIONS:**

- (1) To receive credit you must:
  - (a) **work in a logical fashion, show all your work, indicate your reasoning;**  
no credit will be given for an answer that *just appears*;  
such explanations help with partial credit
  - (b) if a line/box is provided, then:
    - show you work BELOW the line/box
    - put your answer on/in the line/box
  - (c) if no such line/box is provided, then box your answer
- (2) The MARK BOX indicates the problems along with their points.  
Check that your copy of the exam has all of the problems.
- (3) You may **not** use an electronic device, a calculator, books, personal notes.
- (4) During this exam, do not leave your seat unless you have permission. If you have a question, raise your hand. When you finish: turn your exam over, put your pencil down, and raise your hand.
- (5) If you do not make at least 12.5 out of 25 points on Problem 1, then your score for the entire exam will be whatever you made on Problem 1.
- (6) This exam covers (from *Calculus (ET)* by Stewart 6<sup>th</sup> ed.):  
Sections 7.1 – 7.5, 7.8, 11.1 .

**Hints:**

- (1) **You can check your answers to the indefinite integrals by differentiating.**
- (2) **For more partial credit, box your  $u - du$  substitutions.**

**Honor Code Statement**

I understand that it is the responsibility of every member of the Carolina community to uphold and maintain the University of South Carolina's Honor Code.

As a Carolinian, I certify that I have neither given nor received unauthorized aid on this exam.

I understand that if it is determined that I used any unauthorized assistance or otherwise violated the University's Honor Code then I will receive a failing grade for this course and be referred to the academic Dean and the Office of Academic Integrity for additional disciplinary actions.

Furthermore, I have not only read but will also follow the above Instructions.

Signature : \_\_\_\_\_

**1. Fill in the blanks (each worth 1 point).**

**1a.**  $\int \frac{du}{u} = \underline{\hspace{2cm}} |u| + C$

**1b.** If  $a$  is a constant and  $a > 0$  but  $a \neq 1$ , then  $\int a^u du = \underline{\hspace{2cm}} + C$

**1c.**  $\int \cos u du = \underline{\hspace{2cm}} + C$

**1d.**  $\int \sec^2 u du = \underline{\hspace{2cm}} + C$

**1e.**  $\int \sec u \tan u du = \underline{\hspace{2cm}} + C$

**1f.**  $\int \sin u du = \underline{\hspace{2cm}} + C$

**1g.**  $\int \csc^2 u du = \underline{\hspace{2cm}} + C$

**1h.**  $\int \csc u \cot u du = \underline{\hspace{2cm}} + C$

**1i.**  $\int \tan u du = \underline{\hspace{2cm}} + C$

**1j.**  $\int \cot u du = \underline{\hspace{2cm}} + C$

**1k.**  $\int \sec u du = \underline{\hspace{2cm}} + C$

**1l.**  $\int \csc u du = \underline{\hspace{2cm}} + C$

**1m.** If  $a$  is a constant and  $a > 0$  then  $\int \frac{1}{\sqrt{a^2-u^2}} du = \underline{\hspace{2cm}} + C$

**1n.** If  $a$  is a constant and  $a > 0$  then  $\int \frac{1}{a^2+u^2} du = \underline{\hspace{2cm}} + C$

**1o.** If  $a$  is a constant and  $a > 0$  then  $\int \frac{1}{u\sqrt{u^2-a^2}} du = \underline{\hspace{2cm}} + C$

**1p.** Partial Fraction Decomposition. If one wants to integrate  $\frac{f(x)}{g(x)}$  where  $f$  and  $g$  are polynomials and  $[\text{degree of } f] \geq [\text{degree of } g]$ , then one must first do  $\underline{\hspace{2cm}}$

**1q.** Integration by parts formula:  $\int u dv = \underline{\hspace{2cm}}$

**1r.** Trig substitution: (recall that the *integrand* is the function you are integrating)  
if the integrand involves  $a^2-u^2$ , then one makes the substitution  $u = \underline{\hspace{2cm}}$

**1s.** Trig substitution:  
if the integrand involves  $a^2+u^2$ , then one makes the substitution  $u = \underline{\hspace{2cm}}$

**1t.** Trig substitution:  
if the integrand involves  $u^2-a^2$ , then one makes the substitution  $u = \underline{\hspace{2cm}}$

**1u.** trig formula ... your answer should involve trig functions of  $\theta$ , and not of  $2\theta$ :  $\sin(2\theta) = \underline{\hspace{2cm}}$ .

**1v.** trig formula ...  $\cos(2\theta)$  should appear in the numerator:  $\cos^2(\theta) = \frac{\underline{\hspace{2cm}}}{2}$ .

**1w.** trig formula ...  $\cos(2\theta)$  should appear in the numerator:  $\sin^2(\theta) = \frac{\underline{\hspace{2cm}}}{2}$ .

**1x.** trig formula ... since  $\cos^2 \theta + \sin^2 \theta = 1$ , we know that the corresponding relationship between tangent (i.e., tan) and secant (i.e., sec) is  $\underline{\hspace{2cm}}$ .

**1y.**  $\arctan(-\sqrt{3}) = \underline{\hspace{2cm}}$  **RADIANS.** (your answer should be an angle)

2.

$$\int x (5x^2 + 3)^{17} dx =$$

+ C

3.

$$\int \sin^2 x \cos^3 x \, dx =$$

+ C

4.

$$\int e^{5x} \cos(2x) dx = \quad \quad \quad + C$$

Hint: bring to the other side idea.

5.

$$\int x^{\frac{1}{3}} \ln x \, dx =$$

+ C

**6a.** Complete the square by filling in each of the two lines with a (positive or negative) number.

$$x^2 - 6x + 13 = (x + \underline{\hspace{2cm}})^2 + \underline{\hspace{2cm}} .$$

**6b.**

$\int \frac{1}{\sqrt{x^2 - 6x + 13}} dx =$	$+ C$
--	-------

7.

$$\int_1^{\infty} \frac{1}{(3x+1)^4} dx =$$

Warning: write your solution in proper form.

8. Part 8a should help with part 8b.

8a.

$$\int e^{(x^2)} (2x) dx = \qquad + C$$

8b. The functions  $y = e^{x^2}$  and  $y = x^2 e^{x^2}$  do not have elementary antiderivatives.

But the function  $y = (2x^2 + 1) e^{x^2}$  does have an elementary antiderivative.

Evaluate  $\int (2x^2 + 1) e^{x^2} dx$ .

$$\int (2x^2 + 1) e^{x^2} dx = \qquad + C$$