

Math 142, Exam 1, Fall 2009 Solutions

Write your answers as legibly as you can.

There are 11 problems on 6 pages. Problem 1 is worth 10 points. Each of the other problems is worth 9 points. **SHOW** your work. Make your work be coherent and clear. Write in complete sentences whenever this is possible. **CIRCLE** your answer. **CHECK** your answer whenever possible. **No Calculators or Cell phones.**

I will post the solutions on my website a few hours after the exam is finished.

1. **Define the definite integral. Give a complete definition. Be sure to explain all of your notation.**

10 Let $f(x)$ be a function defined on the closed interval $a \leq x \leq b$. For each partition P of the closed interval $[a, b]$ (so, P is $a = x_0 \leq x_1 \leq \dots \leq x_n = b$), let $\Delta_i = x_i - x_{i-1}$, and pick $x_i^* \in [x_{i-1}, x_i]$. The definite integral $\int_a^b f(x)dx$ is the limit over all partitions P as all Δ_i go to zero of $\sum_{i=1}^n f(x_i^*)\Delta_i$.

- 9 2. **State both parts of the Fundamental Theorem of Calculus. Be sure to explain all of your notation.**

Let f be a continuous function defined on the closed interval $[a, b]$.

(a) If $A(x)$ is the function $A(x) = \int_a^x f(t)dt$, for all $x \in [a, b]$, then $A'(x) = f(x)$ for all $x \in [a, b]$.

(b) If $F(x)$ is any antiderivative of $f(x)$, then $\int_a^b f(x)dx = F(b) - F(a)$.

3. **Find $\int 5x \cos(x^2)dx$. Check your answer.**

9 Let $u = x^2$. So, $du = 2xdx$. The original integral is equal to

$$\frac{5}{2} \int \cos u du = \frac{5}{2} \sin u + C = \boxed{\frac{5}{2} \sin(x^2) + C.}$$

Check: The derivative of the proposed answer is

$$\frac{5}{2} 2x \cos(x^2) \checkmark.$$

4. Find $\int \frac{y^2 dy}{\sqrt{3-4y}}$. Check your answer.

We do a change of variables. Suppose that the expression under the radical was just a variable, then we could do algebraic tricks to finish the problem! We make it so! Let $u = 3 - 4y$. It follows that $du = -4dy$. The original problem is equal to

$$\begin{aligned} \int \frac{\left(\frac{3-u}{4}\right)^2 du}{-4\sqrt{u}} &= \frac{1}{-64} \int \frac{(3-u)^2 du}{\sqrt{u}} \\ &= \frac{1}{-64} \int \frac{(9-6u+u^2) du}{\sqrt{u}} = \frac{1}{-64} \int (9u^{-1/2} - 6u^{1/2} + u^{3/2}) du \\ &= \frac{1}{-64} \left(9u^{1/2} \cdot 2 - 6u^{3/2} \cdot \frac{2}{3} + u^{5/2} \cdot \frac{2}{5} \right) + C \\ &= \boxed{\frac{1}{-64} \left(9(3-4y)^{1/2} \cdot 2 - 6(3-4y)^{3/2} \cdot \frac{2}{3} + (3-4y)^{5/2} \cdot \frac{2}{5} \right) + C} \end{aligned}$$

Check: The derivative of the proposed answer is

$$\begin{aligned} &\frac{1}{-64} \left(9(3-4y)^{-1/2}(-4) - 6(3-4y)^{1/2}(-4) + (3-4y)^{3/2}(-4) \right) \\ &= \frac{(3-4y)^{-1/2}(-4)}{-64} (9 - 6(3-4y) + (3-4y)^2) \\ &= \frac{1}{16\sqrt{3-4y}} (9 - 18 + 24y + 9 - 24y + 16y^2) = \frac{1}{16\sqrt{3-4y}} (16y^2) = \frac{y^2}{\sqrt{3-4y}} \checkmark. \end{aligned}$$

5. Find $\int \frac{x}{\sqrt{1-4x^4}} dx$. Check your answer.

We plan to maneuver the given integral into the form

$$\int \frac{du}{\sqrt{1-u^2}} = \arcsin u + C.$$


Let $u = 2x^2$. It follows that $du = 4xdx$. The original problem is equal to

$$\frac{1}{4} \int \frac{du}{\sqrt{1-u^2}} = \frac{1}{4} \arcsin u + C = \boxed{\frac{1}{4} \arcsin(2x^2) + C}.$$

Check: The derivative of the proposed answer is

$$\frac{1}{4} 4x \frac{1}{\sqrt{1-(2x^2)^2}} \checkmark.$$

6. Find the area between $x + y = 0$ and $2 = x + y^2$.



The area is

$$\int_{-1}^2 2 - y^2 - (-y) dy = 2y - \frac{y^3}{3} + \frac{y^2}{2} \Big|_{-1}^2 = \boxed{4 - \frac{8}{3} + 2 - \left(-2 + \frac{1}{3} + \frac{1}{2}\right)}.$$


7. Find the length of the curve parameterized by $x = \cos 2t$, $y = \sin 2t$, with $0 \leq t \leq \pi/2$.

The curve in question is a semi-circle of radius 1; so the length is $\boxed{\pi}$. Also, the length is

9

$$\begin{aligned} \int_0^{\pi/2} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt &= \int_0^{\pi/2} \sqrt{(-2 \sin 2t)^2 + (2 \cos 2t)^2} dt \\ &= \int_0^{\pi/2} \sqrt{4(\sin^2 2t + \cos^2 2t)} dt = \int_0^{\pi/2} \sqrt{4} dt = 2t \Big|_0^{\pi/2} = \boxed{\pi}. \end{aligned}$$

8. Consider the region in the first quadrant bounded by $y = x^2$ and $x = 1$. Revolve this region about $x = 4$. Find the volume of the resulting solid?




Spin the rectangle. Get a shell of volume $2\pi rht$, where $t = dx$, $r = 4 - x$,

4

$h = x^2$. The volume of the solid is:

$$2\pi \int_0^1 (4-x)x^2 dx = 2\pi \int_0^1 (4x^2 - x^3) dx = 2\pi \left(\frac{4x^3}{3} - \frac{x^4}{4} \right) \Big|_0^1 = \boxed{2\pi \left(\frac{4}{3} - \frac{1}{4} \right)}$$


9. Consider the region in the first quadrant bounded by $y = e^x$ and $x = \ln 3$. Revolve this region about the x -axis. Find the volume of the resulting solid.



Spin the rectangle. Get a disk of volume $\pi r^2 t$ where $r = e^x$ and $t = dx$. The volume of the solid is

$$\pi \int_0^{\ln 3} e^{2x} dx = \pi \frac{e^{2x}}{2} \Big|_0^{\ln 3} = \pi \left(\frac{e^{2 \ln 3}}{2} - \frac{1}{2} \right) = \pi \left(\frac{9}{2} - \frac{1}{2} \right) = \boxed{4\pi}$$

10. Consider the solid whose base in the first quadrant of the xy plane is bounded by $y = x^2$ and $y = 1$. Each cross section of this solid perpendicular to the y -axis is a square. Find the volume of the resulting solid.




The slice with y -coordinate y has volume side² t where $t = dy$ and the side is the x -coordinate of the point on $y = x^2$ where the y -coordinate is y . So the

side is \sqrt{y} . The volume of our slice is $\sqrt{y}^2 dy = y dy$. The volume of the solid is

$$\int_0^1 y dy = \frac{y^2}{2} \Big|_0^1 = \boxed{1/2}.$$

11. Suppose that a conical tank is filled with oil which has a density of 50 lb/ft³. The radius at the top of the tank is 5 ft and the tank is 15 ft high. How much work is done in pumping the oil over the edge of the tank?


9



Be sure to draw and label an axis. We have chosen the y -axis to point upward with $y = 0$ being the bottom of the tank. The work to lift the layer of oil with y -coordinate y is

Force \cdot distance = weight \cdot distance = volume \cdot density \cdot distance = $\pi r^2 t \cdot 50 \cdot$ distance,

where $t = dy$, r is computed from similar triangles $\frac{r}{y} = \frac{5}{15}$:



So, $r = \frac{1}{3}y$. The distance to lift the layer of oil whose y -coordinate is y is $15 - y$. The work to lift one layer of oil is:

$$\pi \left(\frac{1}{3}y\right)^2 50(15 - y) dy.$$

The work to lift all of the oil is

$$\begin{aligned}\frac{50\pi}{9} \int_0^{15} (15y^2 - y^3) dy &= \frac{50\pi}{9} \left(\frac{15y^3}{3} - \frac{y^4}{4} \right) \Big|_0^{15} = \frac{50(15)^4\pi}{9} \left(\frac{1}{3} - \frac{1}{4} \right) \\ &= \boxed{\frac{50(15)^4\pi}{9 \cdot 12} \text{ft-lb.}}\end{aligned}$$