

Math 142, Exam 1, Solutions Fall 2012

Write everything on the blank paper provided. **You should KEEP this piece of paper.** If possible: return the problems in order (use as much paper as necessary), use only one side of each piece of paper, and leave 1 square inch in the upper left hand corner for the staple. If you forget some of these requests, don't worry about it – I will still grade your exam.

The exam is worth 50 points. **SHOW** your work. *CIRCLE* your answer. **CHECK** your answer whenever possible.

No Calculators or Cell phones.

The solutions will be posted later today.

1. (7 points) **Define the definite integral. Give a complete definition. Be sure to explain all of your notation.**

Let $f(x)$ be a function defined on the closed interval $a \leq x \leq b$. For each partition P of the closed interval $[a, b]$ (so, P is $a = x_0 \leq x_1 \leq \dots \leq x_n = b$), let $\Delta_i = x_i - x_{i-1}$, and pick $x_i^* \in [x_{i-1}, x_i]$. The definite integral $\int_a^b f(x)dx$ is the limit over all partitions P as all Δ_i go to zero of $\sum_{i=1}^n f(x_i^*)\Delta_i$.

2. (7 points) **State both parts of the Fundamental Theorem of Calculus. Be sure to explain all of your notation.**

Let f be a continuous function defined on the closed interval $[a, b]$.

(a) If $A(x)$ is the function $A(x) = \int_a^x f(t)dt$, for all $x \in [a, b]$, then $A'(x) = f(x)$ for all $x \in [a, b]$.

(b) If $F(x)$ is any antiderivative of $f(x)$, then $\int_a^b f(x)dx = F(b) - F(a)$.

3. (7 points) **Find $\int \frac{e^{\frac{1}{x}}}{x^2} dx$. Check your answer.**

Answer: Let $u = \frac{1}{x}$. Compute $du = -\frac{1}{x^2}dx$. The original integral is equal to

$$= - \int e^u du = -e^u + C = \boxed{-e^{\frac{1}{x}} + C}.$$

Check: The derivative of the proposed answer is

$$-(-1)e^{\frac{1}{x}} \frac{1}{x^2}. \checkmark$$

4. (7 points) **Find** $\int (\ln x)^2 dx$. **Check your answer.**

Answer: Try integration by parts: $\int u dv = uv - \int v du$. Let $u = (\ln x)^2$ and $dv = dx$. It follows that $du = 2 \ln x \left(\frac{1}{x}\right) dx$ and $v = x$. The original integral is

$$x(\ln x)^2 - 2 \int \ln x dx.$$

Use integration by parts again. Let $u = \ln x$ and $dv = dx$. It follows that $du = \left(\frac{1}{x}\right) dx$ and $v = x$. The original integral is

$$x(\ln x)^2 - 2(x \ln x - \int dx) = \boxed{x(\ln x)^2 - 2(x \ln x - x) + C}.$$

Check. The derivative of the proposed answer is

$$\begin{aligned} x \cdot 2 \ln x \left(\frac{1}{x}\right) + (\ln x)^2 - 2 \left(x \left(\frac{1}{x}\right) + \ln x - 1\right) &= 2 \ln x + (\ln x)^2 - 2(1 + \ln x - 1) \\ &= (\ln x)^2. \checkmark \end{aligned}$$

5. (7 points) **Find** $\int \sin 8x \cos 5x dx$. **Check your answer.**

Answer: Add the two identities

$$\begin{aligned} \sin(A + B) &= \sin A \cos B + \cos A \sin B \\ \sin(A - B) &= \sin A \cos B - \cos A \sin B \end{aligned}$$

to get

$$\sin(A + B) + \sin(A - B) = 2 \sin A \cos B.$$

Divide by 2 to get

$$\frac{1}{2} [\sin(A + B) + \sin(A - B)] = \sin A \cos B.$$

Take $A = 8x$ and $B = 5x$. We have

$$\int \sin 8x \cos 5x dx = \frac{1}{2} \int [\sin(13x) + \sin(3x)] dx = \boxed{\frac{-\cos(13x)}{26} - \frac{\cos(3x)}{6} + C}.$$

6. (7 points) **Find** $\int \frac{dx}{\sqrt{x^2 + 16}}$. **Check your answer.**

Answer: Let $x = 4 \tan \theta$. Compute $dx = 4 \sec^2 \theta d\theta$ and

$$x^2 + 16 = 16 \tan^2 \theta + 16 = 16 \sec^2 \theta.$$

The original integral is

$$\begin{aligned} \int \frac{4 \sec^2 \theta d\theta}{4 \sec \theta} &= \int \sec \theta d\theta = \ln |\sec \theta + \tan \theta| + C = \ln \left| \frac{\sqrt{x^2 + 16}}{4} + \frac{x}{4} \right| + C \\ &= \ln \left| \sqrt{x^2 + 16} + x \right| - \ln(4) + C = \boxed{\ln \left| \sqrt{x^2 + 16} + x \right| + K}, \end{aligned}$$

where K is the constant $K = \ln(4) + C$.

Check: The derivative of the proposed answer is

$$\begin{aligned} \frac{\frac{2x}{2\sqrt{x^2+16}} + 1}{\sqrt{x^2+16} + x} &= \frac{\left(\frac{x}{\sqrt{x^2+16}} + 1\right) \sqrt{x^2+16}}{(\sqrt{x^2+16} + x)\sqrt{x^2+16}} = \frac{x + \sqrt{x^2+16}}{(\sqrt{x^2+16} + x)\sqrt{x^2+16}} \\ &= \frac{1}{\sqrt{x^2+16}}. \checkmark \end{aligned}$$

7. (8 points) **Find** $\int \sqrt{5 + 4x - x^2} dx$. **Check your answer.**

Answer: We complete the square: $5 + 4x - x^2 = 5 + 4 - (x^2 - 4x + 4) = 9 - (x - 2)^2$. Let $x - 2 = 3 \sin \theta$. It follows that $9 - (x - 2)^2 = 9 - 9 \sin^2 \theta = 9 \cos^2 \theta$ and $dx = 3 \cos \theta d\theta$. The original problem is equal to

$$\begin{aligned} \int 3 \cos \theta (3 \cos \theta) d\theta &= 9 \int \cos^2 \theta d\theta = \frac{9}{2} \int (1 + \cos 2\theta) d\theta = \frac{9}{2} \left(\theta + \frac{\sin 2\theta}{2} \right) + C \\ &= \frac{9}{2} \left(\theta + \frac{2 \sin \theta \cos \theta}{2} \right) + C = \frac{9}{2} \left(\arcsin \left(\frac{x-2}{3} \right) + \frac{x-2}{3} \frac{\sqrt{5+4x-x^2}}{3} \right) + C \\ &= \boxed{\frac{9}{2} \arcsin \left(\frac{x-2}{3} \right) + \frac{1}{2} (x-2) \sqrt{5+4x-x^2} + C}. \end{aligned}$$

Check. The derivative of the proposed answer is

$$\frac{9}{2} \frac{\frac{1}{3}}{\sqrt{1 - \frac{(x-2)^2}{9}}} + \frac{1}{2} \left(\sqrt{5+4x-x^2} + (x-2) \frac{4-2x}{2\sqrt{5+4x-x^2}} \right)$$

$$\begin{aligned} &= \frac{1}{2} \left(\frac{3}{\frac{1}{3}\sqrt{9-(x-2)^2}} + \sqrt{5+4x-x^2} + (x-2)\frac{2-x}{\sqrt{5+4x-x^2}} \right) \\ &= \frac{1}{2} \left(\frac{9}{\sqrt{5+4x-x^2}} + \sqrt{5+4x-x^2} + \frac{-(x-2)^2}{\sqrt{5+4x-x^2}} \right) \\ &= \frac{1}{2} \left(\frac{9-(x-2)^2}{\sqrt{5+4x-x^2}} + \sqrt{5+4x-x^2} \right) = \frac{1}{2} \left(\sqrt{5+4x-x^2} + \sqrt{5+4x-x^2} \right) \\ &= \sqrt{5+4x-x^2} \checkmark \end{aligned}$$