

Math 544, Exam 2, Fall 2005

Write your answers as legibly as you can on the blank sheets of paper provided. Use only **one side** of each sheet. Be sure to number your pages. Put your solution to problem 1 first, and then your solution to number 2, etc.; although, by using enough paper, you can do the problems in any order that suits you. There are 10 problems. Each problem is worth 5 points. **SHOW** your work. *CIRCLE* your answer. **CHECK** your answer whenever possible. **No Calculators.**

If I know your e-mail address, I will e-mail your grade to you. If I don't already know your e-mail address and you want me to know it, then **send me an e-mail.**

I will post the solutions on my website shortly after the exam is finished.

1. Find the **GENERAL** solution of the system of linear equations $Ax = b$. Also, list three **SPECIFIC** solutions, if possible. **CHECK** that the specific solutions satisfy the equations.

$$A = \begin{bmatrix} 1 & 2 & 2 & 10 \\ 1 & 2 & 3 & 13 \\ 2 & 4 & 5 & 23 \end{bmatrix}, \quad x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix}, \quad b = \begin{bmatrix} -1 \\ -2 \\ -3 \end{bmatrix}.$$

We apply Gaussian Elimination to the augmented matrix

$$\left[\begin{array}{cccc|c} 1 & 2 & 2 & 10 & -1 \\ 1 & 2 & 3 & 13 & -2 \\ 2 & 4 & 5 & 23 & -3 \end{array} \right].$$

Replace $R_2 \mapsto R_2 - R_1$ and $R_3 \mapsto R_3 - 2R_1$ to get

$$\left[\begin{array}{cccc|c} 1 & 2 & 2 & 10 & -1 \\ 0 & 0 & 1 & 3 & -1 \\ 0 & 0 & 1 & 3 & -1 \end{array} \right].$$

Replace $R_1 \mapsto R_1 - 2R_2$ and $R_3 \mapsto R_3 - R_2$ to get

$$\left[\begin{array}{cccc|c} 1 & 2 & 0 & 4 & 1 \\ 0 & 0 & 1 & 3 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right].$$

The general solution is

$$\left\{ \begin{array}{l} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} \in \mathbb{R}^4 \mid \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ -1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -4 \\ 0 \\ -3 \\ 1 \end{bmatrix} \text{ for some numbers} \\ x_2 \text{ and } x_4 \end{array} \right\}.$$

Specific Solution 1. Take $x_2 = x_4 = 0$. We see that $v_1 = \begin{bmatrix} 1 \\ 0 \\ -1 \\ 0 \end{bmatrix}$ is a solution of $Ax = b$ because

$$Av_1 = A \begin{bmatrix} 1 & 2 & 2 & 10 \\ 1 & 2 & 3 & 13 \\ 2 & 4 & 5 & 23 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ -1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1-2 \\ 1-3 \\ 2-5 \end{bmatrix} = \begin{bmatrix} -1 \\ -2 \\ -3 \end{bmatrix} = b. \checkmark$$

Specific Solution 2. Take $x_2 = 1$ and $x_4 = 0$. We see that $v_2 = \begin{bmatrix} -1 \\ 1 \\ -1 \\ 0 \end{bmatrix}$ is a solution of $Ax = b$ because

$$Av_2 = A \begin{bmatrix} 1 & 2 & 2 & 10 \\ 1 & 2 & 3 & 13 \\ 2 & 4 & 5 & 23 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \\ -1 \\ 0 \end{bmatrix} = \begin{bmatrix} -1+2-2 \\ -1+2-3 \\ -2+4-5 \end{bmatrix} = \begin{bmatrix} -1 \\ -2 \\ -3 \end{bmatrix} = b. \checkmark$$

Specific Solution 3. Take $x_2 = 0$ and $x_4 = 1$. We see that $v_3 = \begin{bmatrix} -3 \\ 0 \\ -4 \\ 1 \end{bmatrix}$ is a solution of $Ax = b$ because

$$Av_3 = A \begin{bmatrix} 1 & 2 & 2 & 10 \\ 1 & 2 & 3 & 13 \\ 2 & 4 & 5 & 23 \end{bmatrix} \begin{bmatrix} -3 \\ 0 \\ -4 \\ 1 \end{bmatrix} = \begin{bmatrix} -3-8+10 \\ -3-12+13 \\ -6-20+23 \end{bmatrix} = \begin{bmatrix} -1 \\ -2 \\ -3 \end{bmatrix} = b. \checkmark$$

2. Let U and V be subspaces of \mathbb{R}^n . Does the union $U \cup V$ have to be a subspace of \mathbb{R}^n ? If yes, prove your answer. If no, give a counterexample.

NO. Let $n = 2$, $U = \left\{ \begin{bmatrix} a \\ 0 \end{bmatrix} \mid a \in \mathbb{R} \right\}$, and $V = \left\{ \begin{bmatrix} 0 \\ b \end{bmatrix} \mid b \in \mathbb{R} \right\}$. We see that U and V are subspaces of \mathbb{R}^2 ; but the union $U \cup V$ is not a subspace of \mathbb{R}^2 because this union is not closed under addition. In particular, $u = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ is in U (hence also $U \cup V$) and $v = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ is in V (hence also in $U \cup V$); but the sum $u + v = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ is not in either U or V ; so $u + v \notin U \cup V$.

3. Let $V = \left\{ \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \in \mathbb{R}^3 \mid \begin{array}{l} x_1 + 3x_2 + 4x_3 = 0 \\ 2x_1 + 9x_2 + 5x_3 = 0 \\ 5x_1 + 14x_2 + 41x_3 = 0 \\ -x_1 + 32x_2 + 12x_3 = 0 \end{array} \right\}$. Is V a vector space?

Explain thoroughly.

Yes. The set V is the null space of the matrix

$$\begin{bmatrix} 1 & 3 & 4 \\ 2 & 9 & 5 \\ 5 & 14 & 41 \\ -1 & 32 & 12 \end{bmatrix}.$$

We proved that the null space of every matrix is a vector space.

4. Let $V = \left\{ \begin{bmatrix} x_1 + 3x_2 + 4x_3 \\ 2x_1 + 9x_2 + 5x_3 \\ 5x_1 + 14x_2 + 41x_3 \\ -x_1 + 32x_2 + 12x_3 \end{bmatrix} \in \mathbb{R}^4 \mid x_1, x_2, x_3 \in \mathbb{R} \right\}$. Is V a vector space? **Explain thoroughly.**

Yes. The set V is the column space of the matrix

$$\begin{bmatrix} 1 & 3 & 4 \\ 2 & 9 & 5 \\ 5 & 14 & 41 \\ -1 & 32 & 12 \end{bmatrix}.$$

We proved that the column space of every matrix is a vector space.

5. Let $V = \left\{ \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \in \mathbb{R}^2 \mid x_1 x_2 = 0 \right\}$. Is V a vector space? Explain thoroughly.

NO. The set V is not closed under addition. We see that $u = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $v = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ are in V ; but $u + v = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ is not in V .

6. Define “null space”. Use complete sentences. Include everything that is necessary, but nothing more.

The null space of the matrix A is the set of all column vectors x with $Ax = 0$.

7.

- (a) Define “non-singular”. Use complete sentences. Include everything that is necessary, but nothing more.

The square matrix A is non-singular if the **only** column vector x with $Ax = 0$ is $x = 0$.

(b) Let A be an $n \times n$ matrix. List three statements that are equivalent to the statement “ A is non-singular”.

1. The columns of A are linearly independent.
2. For every vector b in \mathbb{R}^n , the system of equations $Ax = b$ has a unique solution.
3. The matrix A is invertible.

8. Let A and B be 2×2 matrices with A not equal to the zero matrix and $A^2 = AB$. Does A have to equal B ? If yes, prove your answer. If no, give a counterexample.

NO. Let $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 75 & 82 \\ 0 & 0 \end{bmatrix}$. We see that A is not the zero matrix and A does not equal B ; but, A^2 and AB are both equal to $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$.

9. Let A and B be $n \times n$ matrices. At least one of the following statements is always true. Pick a true statement and prove it.
- (a) The column space of A is a subset of the column space of AB .
 - (b) The column space of B is a subset of the column space of AB .

- (c) **The column space of AB is a subset of the column space of A .**
- (d) **The column space of AB is a subset of the column space of B .**

Statement (c) is true. If v is in the column space of AB , then $v = ABw$ for some vector w . Thus, $v = A(Bw)$ for the vector Bw and v is in the column space of A .

10. Let A and B be $n \times n$ matrices. At least one of the following statements is always true. Pick a true statement and prove it.

- (a) **The null space of A is a subset of the null space of AB .**
- (b) **The null space of B is a subset of the null space of AB .**
- (c) **The null space of AB is a subset of the null space of A .**
- (d) **The null space of AB is a subset of the null space of B .**

Statement (b) is true. If v is in the null space of B , then $Bv = 0$. Multiply both sides of the equation on the left by A to see that $ABv = 0$. We now see that v is in the null space of AB .