

PRINT Your Name: \_\_\_\_\_

**Quiz for March 15, 2011**

Let

$$A = \begin{bmatrix} 1 & 2 & 1 & 0 \\ 2 & 5 & 3 & -1 \\ 2 & 2 & 0 & 2 \\ 0 & 1 & 1 & -1 \end{bmatrix}.$$

- (a) Find a basis for the null space of  $A$ .
- (b) Find a basis for the column space of  $A$ .
- (c) Express each column of  $A$  as a linear combination of the vectors in your answer to (b).

**ANSWER:** To find a basis for the null space of  $A$ , we solve  $Ax = 0$ . In other words, we apply Elementary Row Operations to  $A$ . Apply  $R2 \mapsto R2 - 2R1$  and  $R3 \mapsto R3 - 2R1$  to get:

$$\begin{bmatrix} 1 & 2 & 1 & 0 \\ 0 & 1 & 1 & -1 \\ 0 & -2 & -2 & 2 \\ 0 & 1 & 1 & -1 \end{bmatrix}.$$

Apply  $R1 \mapsto R1 - 2R2$ ,  $R3 \mapsto R3 + 2R2$ , and  $R4 \mapsto R4 - R2$  to get

$$\begin{bmatrix} 1 & 0 & -1 & 2 \\ 0 & 1 & 1 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

The solution set of  $Ax = 0$  is the set of all vectors  $x$  with

$$\begin{aligned} x_1 &= x_3 - 2x_4 \\ x_2 &= -x_3 + x_4 \\ x_3 &= x_3 \\ x_4 &= x_4 \end{aligned}$$

In other words the null space of  $A$  is the set of linear combinations of

$$\begin{bmatrix} 1 \\ -1 \\ 1 \\ 0 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} -2 \\ 1 \\ 0 \\ 1 \end{bmatrix}.$$

These two vectors are linearly independent (look at rows 3 and 4); so our answer to (a) is

$$(a) \quad v_1 = \begin{bmatrix} 1 \\ -1 \\ 1 \\ 0 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -2 \\ 1 \\ 0 \\ 1 \end{bmatrix}.$$

The columns in  $A$  that correspond to the leading ones in the reduced matrix are a basis for the column space of  $A$ . Columns 1 and 2 in the reduced matrix have leading ones; so our basis for the column space of  $A$  is columns 1 and 2 of  $A$ ; in other words, our answer to (b) is

$$(b) \quad A_1 = \begin{bmatrix} 1 \\ 2 \\ 2 \\ 0 \end{bmatrix}, \quad A_2 = \begin{bmatrix} 2 \\ 5 \\ 2 \\ 1 \end{bmatrix}.$$

The fact that  $v_1$  is in the null space of  $A$  tells us that  $A_1 - A_2 + A_3 = 0$ ; this tells us how to write the third column of  $A$  in terms of our answer to (b). The fact that  $v_2$  is in the null space of  $A$  tells us that  $-2A_1 + A_2 + A_4 = 0$ . Our answer to (c) is:

$$(c) \quad A_3 = -A_1 + A_2, \quad A_4 = 2A_1 - A_2.$$