

Vaccination Strategies for Epidemic Models

Douglas B. Meade

Department of Mathematics
University of South Carolina
Columbia, SC 29208

~~Phone:~~ (803) 777-6183
~~E-mail:~~ dmeade@math.sc.edu
~~WWW:~~ <http://www.math.sc.edu/~meade/>

27 May 1999

Vaccination Strategies for Epidemic Models

Douglas B. Meade

Department of Mathematics
University of South Carolina
Columbia, SC 29208

Phone: (803) 777-6183
E-mail: meade@math.sc.edu
WWW URL: <http://www.math.sc.edu/~meade/>

27 May 1999

S I R Model

- Assumptions
 - constant population size ($N = S + I + R = 1$)
 - mass-action kinetics for infection
 - exponentially distributed mean time to recovery and natural death
 - no death from infection
 - no vaccination

Ref: Shulgin, Stone, and Agur, *Bull Math Bio* (1998) **60**, 1123–1148.

S I R Model — No Vaccination

$$\dot{S} = \mu - \mu S - \beta SI$$

$$\dot{I} = -\mu I + \beta SI - \gamma I$$

$$\dot{R} = -\mu R + \gamma I$$

$$\dot{N} = 0$$

S I R Model — No Vaccination

- Equilibria

	S	I
infection-free	1	0
endemic	$1/R_0$	$\frac{\mu}{\beta}(R_0 - 1)$

- Reproduction Rate: $R_0 = \frac{\beta}{\mu + \gamma}$
- Stability

	$R_0 < 1$	$R_0 > 1$
infection-free	globally stable	unstable
endemic	unstable	globally stable

S I R Model — Constant Vaccination

- Assumptions

- life-long immunization at birth

$$\begin{aligned}\dot{S} &= (1 - \rho)\mu - \mu S - \beta SI \\ \dot{I} &= -\mu I + \beta SI - \gamma I \\ \dot{R} &= \rho\mu - \mu R + \gamma I \\ \dot{N} &= 0\end{aligned}$$

S I R Model — Constant Vaccination

- Equilibria

	S	I
infection-free	$1 - \rho$	0
endemic	$1/R_0$	$\frac{\mu}{\beta}(R_0 - 1) - \frac{\mu}{\mu + \gamma}\rho$

- Reproduction Rate: $R_\rho = (1 - \rho) R_0$
- Critical Vaccination Rate: $\rho_0 = 1 - \frac{1}{R_0}$
- Stability

	$R_\rho < 1$ ($\rho > \rho_0$)	$R_\rho > 1$ ($\rho < \rho_0$)
infection-free	globally stable	unstable
endemic	unstable	globally stable

S I R Model — Pulse Vaccination

- Assumptions

- fixed portion of susceptibles vaccinated every T years

$$\dot{S} = \mu - \mu S - \beta SI - \sum_{n=0}^{\infty} \rho S(nT^-) \delta(t - nT)$$

$$\dot{I} = -\mu I + \beta SI - \gamma I$$

$$\dot{R} = -\mu R + \gamma I + \sum_{n=0}^{\infty} \rho S(nT^-) \delta(t - nT)$$

$$\dot{N} = 0$$

S I R Model — Pulse Vaccination

- Infection-Free Equilibrium

– initial value problem:

$$\dot{S} = \mu - \mu S - \sum_{n=0}^{\infty} \rho S(nT^-) \delta(t - nT), \quad S(0) = S_0$$

– solution:

$$S(t) = \begin{cases} S^{(n)}(t), & t \in [nT, (n+1)T) \\ S_{n+1}, & t = (n+1)T \end{cases}$$

$$S^{(n)}(t) = 1 - (1 - S_n) e^{-\mu(t-nT)}$$

$$S_{n+1} = S((n+1)T^+)$$

$$= (1 - \rho) S^{(n)}((n+1)T^-)$$

S I R Model — Pulse Vaccination

– construction of periodic equilibrium solution:

* if $S_\infty = \lim_{n \rightarrow \infty} S_n$ exists, then S_∞ is a fixed point of

$$F(x) = (1 - \rho)(1 - (1 - x)e^{-\mu T})$$

* unique fixed point is locally stable

$$S_\infty = \frac{(1 - \rho)(e^{\mu T} - 1)}{e^{\mu T} - (1 - \rho)}$$

* periodic equilibrium solution:

$$S^{(\infty)}(t) = 1 - (1 - S_\infty)e^{-\mu(t \bmod T)}$$

S I R Model — Pulse Vaccination

- local stability:

$$\frac{1}{T} \int_0^T S \, dt < S_c$$

where $S_c = 1/R_0$ is the endemic equilibrium with no vaccination

- maximum vaccination period:

$$\frac{(\mu T - \rho)(e^{\mu T} - 1) + \mu \rho T}{\mu T(e^{\mu T} - (1 - \rho))} = \frac{\mu + \gamma}{\beta}$$

Application: Rubella (Measles)

- Parameters

mean infectious period	3.5 days	$\gamma = 100$
mean lifespan	50 years	$\mu = 0.02$
contact rate		$\beta = 1800$

- No Vaccination:
 - $R_0 \approx 18 \implies$ endemic state is stable
 - endemic equilibrium: $S \approx 0.05$
- Constant Vaccination:
 - $\rho_0 \approx 0.95$
 - reasonable for school age, but not pre-school (35-50%)

S E I R Model

- Assumptions
 - exposed period prior to becoming infectious
 - non-constant population size ($N = S + E + I + R$)
 - * immigration
 - * birth
 - * death from infection
 - mass-action kinetics for infection
 - exponentially distributed mean time to recovery and natural death
 - no vaccination

S E I R Model — No Vaccination

$$\dot{S} = \Lambda + \mu_0 N - \mu_1 S - \beta SI$$

$$\dot{E} = -\mu_1 E + \beta SI - \delta E$$

$$\dot{I} = -\mu_1 I + \delta E - \gamma I - \epsilon I$$

$$\dot{R} = -\mu_1 R + \gamma I$$

$$\dot{N} = \Lambda - (\mu_1 - \mu_0)N - \epsilon I$$

Note:

- all parameters are positive
- to prevent exponential growth in total population, assume death-dominant ($\mu_1 > \mu_0$)
- S I R model obtained with $\epsilon = 0$, $\delta = \infty$, $\frac{\Lambda}{\mu_1 - \mu_0} = 1$

S E I R Model — No Vaccination

- Equilibria

	infection-free	endemic
S	$\frac{\Lambda}{\mu_1 - \mu_0}$	$\frac{(\mu_1 + \delta)(\mu_1 + \gamma + \epsilon)}{\beta\delta}$
E	0	$\frac{\mu_1(\mu_1 + \gamma + \epsilon)}{\beta\delta} \frac{A_0}{B_0}$
I	0	$\frac{\mu_1}{\beta} \frac{A_0}{B_0}$
R	0	$\frac{\gamma}{\beta} \frac{A_0}{B_0}$
N	$\frac{\Lambda}{\mu_1 - \mu_0}$	$\frac{(\beta\Lambda + \epsilon\mu_1)(\mu_1 + \delta)(\mu_1 + \gamma + \epsilon) - \epsilon\delta\beta\Lambda}{\beta B_0}$

where

$$A_0 = \beta\delta\Lambda - (\mu_1 - \mu_0)(\mu_1 + \delta)(\mu_1 + \gamma + \epsilon)$$

$$B_0 = \epsilon\delta\Lambda + (\mu_1 - \mu_0)(\mu_1 + \delta)(\mu_1 + \gamma + \epsilon)$$

S E I R Model — No Vaccination

- Reproduction Rate: $R_0 = \frac{\Lambda}{\mu_1 - \mu_0} \frac{\delta}{\mu_1 + \delta} \frac{\beta}{\mu_1 + \gamma + \epsilon}$
- Stability

	$R_0 < 1$	$R_0 > 1$
infection-free	globally stable	unstable
endemic	unstable	globally stable

S E I R Model — Constant Vaccination

- Assumptions

- life-long immunization at birth

$$\dot{S} = \Lambda + (1 - \rho)\mu_0 N - \mu_1 S - \beta SI$$

$$\dot{E} = -\mu_1 E + \beta SI - \delta E$$

$$\dot{I} = -\mu_1 I + \delta E - \gamma I - \epsilon I$$

$$\dot{R} = \rho\mu_0 N - \mu_1 R + \gamma I$$

$$\dot{N} = \Lambda - (\mu_1 - \mu_0)N - \epsilon I$$

S E I R Model — Constant Vaccination

- Equilibria

	infection-free	endemic
S	$\frac{\Lambda}{\mu_1 - \mu_0} \left(1 - \frac{\mu_0}{\mu_1} \right)$	$\frac{(\mu_1 + \delta)(\mu_1 + \gamma + \epsilon)}{\beta \delta}$
E	0	$\frac{\mu_1(\mu_1 + \gamma + \epsilon)}{\beta \delta} \frac{A_\rho}{B_\rho}$
I	0	$\frac{\mu_1}{\beta} \frac{A_\rho}{B_\rho}$
R	$\frac{\Lambda}{\mu_1 - \mu_0} \frac{\mu_0}{\mu_1}$	$\frac{\gamma}{\beta} \frac{A_\rho}{B_\rho} + \frac{\mu_0}{\mu_1} \rho N$
N	$\frac{\Lambda}{\mu_1 - \mu_0}$	$\frac{(\beta \Lambda + \epsilon \mu_1)(\mu_1 + \delta)(\mu_1 + \gamma + \epsilon) - \epsilon \delta \beta \Lambda}{\beta B_\rho}$

where

$$A_\rho = A_0 - \frac{\mu_0}{\mu_1} \rho \beta \delta \Lambda, \quad B_\rho = B_0 + \frac{\mu_0}{\mu_1} \rho \epsilon \delta \mu_0$$

S E I R Model — Constant Vaccination

- Reproduction Rate: $R_\rho = \left(1 - \frac{\mu_0}{\mu_1} \rho\right) R_0$
- Critical Vaccination Rate: $\rho_0 = \frac{\mu_1}{\mu_0} \left(1 - \frac{1}{R_0}\right)$
- Stability

	$R_\rho < 1$ ($\rho > \rho_0$)	$R_\rho > 1$ ($\rho < \rho_0$)
infection-free	globally stable	unstable
endemic	unstable	globally stable

S E I R Model — Pulse Vaccination

- Assumptions

– fixed portion of susceptibles vaccinated every T years

$$\dot{S} = \Lambda + \mu_0 N - \mu_1 S - \beta SI$$

$$- \sum_{n=0}^{\infty} \rho S(nT^-) \delta(t - nT)$$

$$\dot{E} = -\mu_1 E + \beta SI - \delta E$$

$$\dot{I} = -\mu_1 I + \delta E - \gamma I - \epsilon I$$

$$\dot{R} = -\mu_1 R + \gamma I$$

$$+ \sum_{n=0}^{\infty} \rho S(nT^-) \delta(t - nT)$$

$$\dot{N} = \Lambda - (\mu_1 - \mu_0)N - \epsilon I$$

S E I R Model — Pulse Vaccination

- Infection-Free Equilibrium

– initial value problem:

$$\begin{aligned} \dot{S} &= \Lambda + \mu_0 N - \mu_1 S - \sum_{n=0}^{\infty} \rho_S(nT^-) \delta(t - nT) \\ \dot{R} &= -\mu_1 R + \sum_{n=0}^{\infty} \rho_S(nT^-) \delta(t - nT) \\ \dot{N} &= \Lambda - (\mu_1 - \mu_0) N \end{aligned}$$

S E I R Model — Pulse Vaccination

– solution:

$$S(t) = \begin{cases} S^{(n)}(t), & t \in [nT, (n+1)T) \\ S_{n+1}, & t = (n+1)T \end{cases}$$

where, for all $n=0, 1, 2, \dots$,

$$\begin{aligned} S^{(n)}(t) &= \frac{\Lambda}{\mu_1 - \mu_0} + \left(S_n - \frac{\Lambda}{\mu_1 - \mu_0} \right) e^{-\mu_1(t-nT)} \\ &\quad + \left(N_0 - \frac{\Lambda}{\mu_1 - \mu_0} \right) \left(1 - e^{-\mu_0(t-nT)} \right) e^{-(\mu_1 - \mu_0)t} \\ S_{n+1} &= S((n+1)T^+) \\ &= (1 - \rho) S^{(n)}((n+1)T^-) \end{aligned}$$

S E I R Model — Pulse Vaccination

— construction of periodic equilibrium solution:

* if $S_\infty = \lim_{n \rightarrow \infty} S_n$ exists, then S_∞ is a fixed point of

$$F(x) = (1 - \rho) \left(\frac{\Lambda}{\mu_1 - \mu_0} + \left(x - \frac{\Lambda}{\mu_1 - \mu_0} \right) e^{-\mu_1 T} \right)$$

* unique fixed point is locally stable

$$S_\infty = \frac{\Lambda}{\mu_1 - \mu_0} \frac{(1 - \rho)(e^{\mu T} - 1)}{e^{\mu T} - (1 - \rho)}$$

S E I R Model — Pulse Vaccination

* asymptotically periodic equilibrium solution:

$$S^{(\infty)}(t) = \frac{\Lambda}{\mu_1 - \mu_0} + \left(S_\infty - \frac{\Lambda}{\mu_1 - \mu_0} \right) e^{-\mu_1 (t \bmod T)}$$

$$+ \left(N_0 - \frac{\Lambda}{\mu_1 - \mu_0} \right) \left(1 - e^{-\mu_0 (t \bmod T)} \right)$$

$$\times e^{-(\mu_1 - \mu_0)t}$$

— ...

Application: Rubella (Measles)

- Parameters

mean exposed period	9 days	$\delta = 40$
mean infectious period	6 days	$\gamma = 60$
mean lifespan	80 years	$\mu = 0.0125$
contact rate	“high”	$\beta =$
immigration rate	???	$\Lambda =$
net birth rate	???	$\mu_0 =$
net death rate	???	$\mu_1 =$