

SCCC 411B Exam 2 Spring, 1997 Name: \_\_\_\_\_

**Instructions.** There are 100 points. Answer all questions. Be sure to supply adequate explanation for your answers.

1. (35 points) The Leslie matrix for a population of New Zealand sheep is given by the following data, where  $F_i$  denotes the fecundity and  $P_i$  the probability of survival to the next age class.

age (yr)	0	1	2	3	4	5	6	7	8	9	10	11
$F_i$	.0	.04	.39	.47	.48	.55	.54	.50	.47	.46	.43	.42
$P_i$	.84	.98	.96	.95	.93	.90	.85	.79	.69	.56	.37	.0

- a. Write out the upper left  $4 \times 4$  corner of the matrix  $L$ .
- b. If the current population consists of 100 two year olds and 100 three year olds how many individuals will be in each age class one year from now?
- c. The dominant eigenvalue for  $L$  is  $\lambda = 1.18$ , and the corresponding eigenvector is  $[\.24, \.17, \.14, \.12, \.10, \.08, \.06, \.04, \.03, \.02, \.01, \.0]$ . The dominant eigenvalue for  $L^T$  is  $\mu = 1.18$ , and the corresponding eigenvector is  $[1.0, 1.4, 1.6, 1.6, 1.5, 1.3, 1.1, 0.9, 0.8, 0.6, 0.5, 0.4]$ . Over the long term does the population grow, decline, or remain stable? Why? In the long term what percent of the population is 7 years old or older? Which age class has the highest reproductive value? Is this the age class with the highest fecundity? If not, explain the apparent discrepancy.

- d. Recall that the elasticity  $e_{ij}$  times percent change in  $L_{ij}$  gives percent change in  $\lambda$ . We have calculated that  $e_{13} = 0.037$  and  $e_{18} = 0.014$ . Suppose the fecundity of the two year olds rises from .39 to .44 and the fecundity of the seven year olds drops from .50 to .40. Which has a larger effect on  $\lambda$  in terms of percent change? How do you account for this intuitively?

2. (20 points)

- a. You are given the plot  $N_{t+1} = F(N_t)$ . Based on this graph, what is the steady state  $\bar{N}$ ? If  $N_0 = 5$ , use the graph to compute  $N_4$ .

- b. You are given the plots  $N_{t+1} = G(N_t)$ ,  $N_{t+2} = G(N_{t+1}) = G(G(N_t))$ , and the line  $N_{t+2} = N_t$ . Interpret the points A, B, and C in terms of



- c. What fraction of  $N_t$  is available for reproduction, either by hiding, or by being exposed but lucky?

- d. Explain how the equations given below reflect our description of the model.

$$N_{t+1} = \lambda N_t \left( \frac{bK}{N_t} + \left( 1 - \frac{bK}{N_t} \right) e^{-aP_t} \right)$$

$$P_{t+1} = cN_t \left( 1 - \frac{bK}{N_t} \right) (1 - e^{-aP_t})$$

- e. There is a potential flaw in this model. What does it predict for very low host populations (for example, around  $\frac{1}{2}bK$ )?

- f. With the parameter values  $a = 0.2$ ,  $b = 0.1$ ,  $c = 1$ ,  $K = 100$ ,  $\lambda = 2.8$ , we find that there is a steady state  $\bar{N} = 29.25$ ,  $\bar{P} = 18.81$ ; the corresponding Jacobian matrix has eigenvalues  $0.08 \pm 0.49i$ . Describe the behavior of the system near the equilibrium. In what respect is this result a bit surprising? Upon reflection how do you account for this?

4. (15 points) Give brief definitions of the following terms. Illustrative pictures might also be appropriate.
- a. non-linear difference equation

b. Lefkowitz matrix

c. memoryless random process